

PRODUCT INFORMATION SHEET

Mouse Monoclonal Antibody to SRA

Format: PURE RUO REF IQP-1064P Volume: 0.1ml

Description

Clone 1D4H8

Isotype Mouse IgG1

Specificity SRA

Alternative names

SRAP; STRAA1

Species Human

Immunogen Purified recombinant fragment of SRA expressed in E. Coli.

Μw

Format Ascitic fluid containing 0.03% sodium azide.

Summary

Steroid receptor RNA activator 1 (SRA), with 237-amino acid protein (about 27kDa), belongs to the growing family of functional non-coding RNAs. SRA was originally described as the first functional noncoding RNA able to specifically coactivate the activity of steroid receptors. Specifically, SRA exists as both an RNA transcript that forms a complex with steroid receptor coactivator-1 and as a stably expressed protein. Its expression is strongly up-regulated in many human tumors of the breast, uterus, and ovary, suggesting a potential role in pathogenesis. Although coactivation of steroid-dependent transcription by SRA is accompanied by a proliferative response, overexpression is not in itself sufficient to induce turmorigenesis.

Applications

Western Bloting: 1/500 - 1/2000.Immunohistochemistry: 1/200 - 1/1000.ELISA: Propose dilution 1/10000.Not yet tested in other applications.Determining optimal working dilutions by titration test.

Limitations

- 1. Conjugates with brighter fluorochromes, like PE and APC, will have a greater separation than those with dyes like FITC and CyQ. When populations overlap, the percentage of positive cells using a selected marker can be affected by the choice of fluorescent label.
- Use of monoclonal antibodies in patient treatment can interfere with antigen target recognition by this reagent. This should be taken into account when samples are analyzed from patients treated in this fashion. IQ Products has not characterized the effect of the presence of therapeutic antibodies on the performance of this reagent.
- 3. Reagents can be used in different combinations, therefore laboratories need to become familiar performance characteristics of each antibody in relation with the combined markers in normal and abnormal samples.
- 4. Reagent data performance is based on EDTA-treated blood. Reagent performance can be affected by the use of other anticoagulants.

IQP-1064 – SRA Version 1

▲ ♦ 4 8

Handling and Storage

Antibodies are supplied in 0.01 M sodium phosphate, 0.15 M NaCl; pH 7.3, 0.03% sodiumazide (NaN3) or as ascetic fluid containing 0.03% sodiumazide. Store the vials at 2-8 °C for a maximum of 2 weeks and store at -20°C for longer term storage. Monoclonal antibodies should be protected from prolonged exposure to light when conjugated with fluorochromes. Reagents are stable for the period shown on the vial label when stored properly.

Warranty

Products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied, which extend beyond the description on the label of the product. IQ Products is not liable for property damage, personal injury, or economic loss caused by the product.

Characterization

To ensure consistently high-quality reagents, each batch of monoclonal antibody is tested for conformance with characteristics of a standard reagent. Representative flow cytometric data is included in this data sheet.

Warning

All products contain sodiumazide. This chemical is poisonous and hazardous. Handling should be done by trained staff only.

References 1. Rainer B. Lanz, Steven S. Chua, Niall Barron. Mol. Cell. Biol, Oct 2003; 23: 7163 - 7176. 2. Shilpa Chooniedass-Kothari, Mohammad Kariminia Hamedani, Sandy Troup. Int J Cancer. 2006 Feb 15;118(4):1054-9

3. S. Chooniedass-Kothari, E. Emberley, M. K.

Explanation of used symbols

Consult instructions for use REF Catalogue number V Sufficient for \triangle Caution, consult accompanying document 巻 Keep away from (sun)light 8 Biological risks Temperature limitation (°C) right fluorescence RUO For Research Use Only LOT Batch code Use by yyyy-mm-dd Manufacturer Authorized Representative in the European Community

EC REP Conformité Européenne (European Conformity)

P F R C	PURE FITC R-PE CyQ APC	Label - tandem purified material FITC PE PE-Cy5.18	Ex -max (nm) - 488 488, 532 488, 532 595, 633, 635, 647	Em -max (nm) - 519 578 667 660
PC	PerCP		488, 532	678
PCC	PerCP-Cy5.5		488, 532	695

IQ Products BV

Rozenburglaan 13a 9727 DL Groningen, The Netherlands

+31 (0)50 57 57 000 +31 (0)50 57 57 002 ļ

marketing@igproducts.nl S Technical Orders orders@igproducts.nl

www.iqproducts.nl

IQP-1064 - SRA Version 1