

PRODUCT INFORMATION SHEET

Mouse Monoclonal Antibody to Tyk2

Format: PURE RUO REF IQP-1080P Volume: 0.1ml

Description \prod i

8G8E9, 8G9H5,8G8B3 Clone

Isotype Mouse IgG1

Specificity Tyk2 **Alternative** JTK1 names **Species** Human

Immunogen Purified recombinant fragment of Tyk2 expressed in E. Coli.

Mw

Format Ascitic fluid containing 0.03% sodium azide.

Summary

Tyk2 (tyrosine kinase 2), with 1187-amino acid protein (about 131kDa), belongs to the family of non-receptor janus tyrosine kinases, which also includes Jak1, Jak2, and Jak3. Kinases of the Jak family regulate a spectrum of cellular functions downstream of activated cytokine receptors in the lympho-hematopoietic system. Tyk2 is activated by a variety of cytokines: IFN-alpha, IFN-beta, IL-6, IL-10, IL-12, and IL-13 and promotes IFN-gamma production by Th1-type CD4 cells. Tyk2 can be viewed as a dual-function Jak, mediating both pro-inflammatory and anti-inflammatory cytokine responses. Tyk2 is also an important regulator of lymphoid tumor surveillance.

Applications Western Bloting: 1/500 - 1/2000.ELISA: Propose dilution 1/10000.Not yet tested in other applications. Determining optimal working dilutions by titration test.

Limitations

- 1. Conjugates with brighter fluorochromes, like PE and APC, will have a greater separation than those with dyes like FITC and CyQ. When populations overlap, the percentage of positive cells using a selected marker can be affected by the choice of fluorescent label.
- 2. Use of monoclonal antibodies in patient treatment can interfere with antigen target recognition by this reagent. This should be taken into account when samples are analyzed from patients treated in this fashion. IQ Products has not characterized the effect of the presence of therapeutic antibodies on the performance of this reagent.
- Reagents can be used in different combinations, therefore laboratories need to become familiar performance characteristics of each antibody in relation with the combined markers in normal and abnormal samples.
- 4. Reagent data performance is based on EDTA-treated blood. Reagent performance can be affected by the use of other anticoagulants.

IQP-1080 - Tyk2 Version 1

△ ♦ / * □

Handling and Storage

Antibodies are supplied in 0.01 M sodium phosphate, 0.15 M NaCl; pH 7.3, 0.03% sodiumazide (NaN3) or as ascetic fluid containing 0.03% sodiumazide. Store the vials at 2-8 °C for a maximum of 2 weeks and store at -20°C for longer term storage. Monoclonal antibodies should be protected from prolonged exposure to light when conjugated with fluorochromes. Reagents are stable for the period shown on the vial label when stored properly.

Warranty

Products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied, which extend beyond the description on the label of the product. IQ Products is not liable for property damage, personal injury, or economic loss caused by the product.

Characterization

To ensure consistently high-quality reagents, each batch of monoclonal antibody is tested for conformance with characteristics of a standard reagent. Representative flow cytometric data is included in this data sheet.

Warning

All products contain sodiumazide. This chemical is poisonous and hazardous. Handling should be done by trained staff only.

References 1. Michael H. Shaw, Gordon J. Freeman, Mark F. Scott. J. Immunol., Jun 2006; 176: 7263-7271. 2. Yohei Seto, Hiroshi Nakajima, Akira Suto. J. Immunol., Jan 2003; 170: 1077.

Explanation of used symbols

Consult instructions for use

REF
Sufficient for
Caution, consult accompanying document
Keep away from (sun)light
Biological risks

Temperature limitation (°C)

For Research Use Only

Batch code

Use by yyyy-mm-dd Manufacturer

Authorized Representative in the European Community Conformité Européenne (European Conformity)

P F R C A	PURE FITC R-PE CyQ APC PerCP	Label - tandem purified material FITC PE PE-Cy5.18	Ex -max (nm) - 488 488, 532 488, 532 595, 633, 635, 647 488, 532	Em -max (nm) - 519 578 667 660 678
PCC	PerCP-Cy5.5		488, 532	695



LOT

EC REP

IQ Products BV

Rozenburglaan 13a 9727 DL Groningen, The Netherlands

** +31 (0)50 57 57 000 +31 (0)50 57 57 002

Technical marketing@iqproducts.nl
 Orders orders@iqproducts.nl

<u>www.iqproducts.nl</u>

IQP-1080 - Tyk2 Version 1