

PRODUCT INFORMATION SHEET

Mouse Monoclonal Antibody to PR						
	Format: PURE RUO REF IQP-1093P Volume: 0.1ml					
	Description					
Clone	2F12B4,2F12H2					
Isotype	Mouse IgG1					
Specificity	PR					
Alternative names	PR; NR3C3; PGR					
Species	Human					
[mmunogen	Purified recombinant fragment of PR expressed in E. Coli.					
Mw						
Format	Ascitic fluid containing 0.03% sodium azide.					
Summary	PR(progesterone receptor), with 933-amino acid protein (about 110kDa), a member of the steroid receptor superfamily, mediates the physiologic effects of progesterone, PR is mediated by two functionally different isoforms of the progesterone receptor, the full length PR-B and the short form PR-A. The PR-A and PR-B proteins are 94 kDa and 114 kDa respectively. That are equimolar in the normal breast but dysregulated in advanced disease. PR is prognostic markers in breast cancers irrespective of the patient's progestational status Human progesterone.					
Applications	Western Bloting: 1/500 - 1/2000.Immunohistochemistry: 1/200 - 1/1000.ELISA: Propose dilution 1/10000.Not yet tested in other applications.Determining optimal working dilutions by titration test.					
Limitations	 Conjugates with brighter fluorochromes, like PE and APC, will have a greater separation than those with dyes like FITC and CyQ. When populations overlap, the percentage of positive cel 					
	using a selected marker can be affected by the choice of fluorescent label.Use of monoclonal antibodies in patient treatment can interfere with antigen target recogniti by this reagent. This should be taken into account when samples are analyzed from patients					
	 treated in this fashion. IQ Products has not characterized the effect of the presence of the rapeutic antibodies on the performance of this reagent. Reagents can be used in different combinations, therefore laboratories need to become familiar performance characteristics of each antibody in relation with the combined markers normal and abnormal samples. 					
	 Reagent data performance is based on EDTA-treated blood. Reagent performance can be affected by the use of other anticoagulants. 					

⚠ 💩 🔏 🛣 🎴

Handling and Storage

- Antibodies are supplied in 0.01 M sodium phosphate, 0.15 M NaCl; pH 7.3, 0.03% sodiumazide (NaN3) or as ascetic fluid containing 0.03% sodiumazide. Store the vials at 2-8 °C for a maximum of 2 weeks and store at -20°C for longer term storage. Monoclonal antibodies should be protected from prolonged exposure to light when conjugated with fluorochromes. Reagents are stable for the period shown on the vial label when stored properly.
- **Warranty** Products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied, which extend beyond the description on the label of the product. IQ Products is not liable for property damage, personal injury, or economic loss caused by the product.

Characterization

- To ensure consistently high-quality reagents, each batch of monoclonal antibody is tested for conformance with characteristics of a standard reagent. Representative flow cytometric data is included in this data sheet.
- **Warning** All products contain sodiumazide. This chemical is poisonous and hazardous. Handling should be done by trained staff only.

References 1. McGuire, W.L., et al. 1986. Cancer.Surv.5:527-536.

- 2. Helena CD, Curt WB, Paul V, et al. 2000. J.Clin. Pathol. 53:201-205.
- 3. Wen, D.X., et al. 1994.Mol.Cell.Biol.14:8356-8364.
- 4. GM Clark, CK Osborne, WL McGuire.1984.J. Clin. Oncol., 2:1102-1109.

Explanation of used symbols

REF Ø	Consult instructions for use Catalogue number Sufficient for Caution, consult accompanying document				
<u>*</u> ⊗	Keep away from (sun)light Biological risks				
KUO LOT	Temperature limitation (°C) For Research Use Only				
	Batch code Dingine inconcessee Concession Co				
EC REP	Authorized Representative in the European Community Conformité Européenne (European Conformity)				

		Label - tandem	Ex -max (nm)	Em -max (nm)
Р	PURE	purified material	-	-
F	FITC	FITC	488	519
R	R-PE	PE	488, 532	578
С	CyQ	PE-Cy5.18	488, 532	667
Α	APC		595, 633, 635, 647	660
PC	PerCP		488, 532	678
PCC	PerCP-Cy5.5		488, 532	695

IQ Products BV

Rozenburglaan 13a

9727 DL Groningen, The Netherlands

- +31 (0)50 57 57 000
- +31 (0)50 57 57 002
- Technical <u>marketing@iqproducts.nl</u>
- Orders <u>orders@iqproducts.nl</u>
- <u>www.iqproducts.nl</u>