

# PRODUCT INFORMATION SHEET

# **Mouse Monoclonal Antibody to TrkA**

Format: PURE RUO REF IQP-1283P Volume: 0.1ml

**Description**  $\prod$ i

Clone 6B2

Mouse IgG1 **Isotype** 

**Specificity** TrkA

**Alternative** names

NTRK1, MTC, TRK, TRK1

**Species** Human

Purified recombinant extracellular fragment of human TrkA (aa33-423) fused with hIgGFc tag Immunogen

expressed in HEK293 cell line.

**Format** Ascitic fluid containing 0.03% sodium azide.

**Summary** 

TrkA, also known as NTRK1, MTC, TRK, TRK1. It is a member of the neurotrophic tyrosine kinase receptor (NTKR) family. This kinase is a membrane-bound receptor that, upon neurotrophin binding, phosphorylates itself and members of the MAPK pathway. The presence of this kinase leads to cell differentiation and may play a role in specifying sensory neuron subtypes. Mutations in this gene have been associated with congenital insensitivity to pain, anhidrosis, self-mutilating behavior, mental retardation and cancer. Alternate transcriptional splice variants of this gene have been found, but only three have been characterized to date.

Applications Western Bloting: 1/500 - 1/2000.Immunofluorescence: 1/200 - 1/1000.ELISA: Propose dilution 1/10000.Not yet tested in other applications. Determining optimal working dilutions by titration

#### Limitations

- 1. Conjugates with brighter fluorochromes, like PE and APC, will have a greater separation than those with dyes like FITC and CyQ. When populations overlap, the percentage of positive cells using a selected marker can be affected by the choice of fluorescent label.
- 2. Use of monoclonal antibodies in patient treatment can interfere with antigen target recognition by this reagent. This should be taken into account when samples are analyzed from patients treated in this fashion. IQ Products has not characterized the effect of the presence of therapeutic antibodies on the performance of this reagent.
- 3. Reagents can be used in different combinations, therefore laboratories need to become familiar performance characteristics of each antibody in relation with the combined markers in normal and abnormal samples.
- 4. Reagent data performance is based on EDTA-treated blood. Reagent performance can be affected by the use of other anticoagulants.

IOP-1283 - TrkA Version 1

# **△ ♦ / \*** □

### **Handling and Storage**

Antibodies are supplied in 0.01 M sodium phosphate, 0.15 M NaCl; pH 7.3, 0.03% sodiumazide (NaN3) or as ascetic fluid containing 0.03% sodiumazide. Store the vials at 2-8 °C for a maximum of 2 weeks and store at -20°C for longer term storage. Monoclonal antibodies should be protected from prolonged exposure to light when conjugated with fluorochromes. Reagents are stable for the period shown on the vial label when stored properly.

#### Warranty

Products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied, which extend beyond the description on the label of the product. IQ Products is not liable for property damage, personal injury, or economic loss caused by the product.

### Characterization

To ensure consistently high-quality reagents, each batch of monoclonal antibody is tested for conformance with characteristics of a standard reagent. Representative flow cytometric data is included in this data sheet.

#### Warning

All products contain sodiumazide. This chemical is poisonous and hazardous. Handling should be done by trained staff only.

**References** 1. DNA Repair (Amst). 2008 Oct 1;7(10):1757-64. 2. Traffic. 2008 Jul;9(7):1146-56.

#### **Explanation of used symbols**

Consult instructions for use

REF

Catalogue number

Sufficient for

Caution, consult accompanying document

Keep away from (sun)light

Biological risks
Temperature limitation (°C)

RUO For Research Use Only
LOT Batch code

Use by yyyy-mm-dd Manufacturer

Authorized Representative in the European Community Conformité Européenne (European Conformity)

P F R C A	PURE FITC R-PE CyQ APC PerCP	Label - tandem purified material FITC PE PE-Cy5.18	Ex -max (nm) - 488 488, 532 488, 532 595, 633, 635, 647 488, 532	Em -max (nm) - 519 578 667 660 678
PCC	PerCP-Cy5.5		488, 532	695



EC REP

IQ Products BV

Rozenburglaan 13a 9727 DL Groningen, The Netherlands

\*\* +31 (0)50 57 57 000 \*\* +31 (0)50 57 57 002

✓ Technical marketing@iqproducts.nl✓ Orders orders@iqproducts.nl

<u>www.iqproducts.nl</u>

IOP-1283 - TrkA Version 1