

PRODUCT INFORMATION SHEET

Mouse Monoclonal Antibody to CD276

Format: PURE RUO REF IQP-1469P Volume: 0.1ml

Description \prod i

Clone 6A1

Mouse IgG1 **Isotype Specificity** CD276

Alternative names

B7H3; B7-H3; 4Iq-B7-H3; CD276

Species Human

Immunogen Purified recombinant fragment of human CD276 expressed in E. Coli.

Mw

Format

Ascitic fluid containing 0.03% sodium azide.

Summary

Costimulatory B7 molecules (e.g., B7-1, or CD80; MIM 112203) signal through CD28 (MIM 186760) family molecules such as CD28, CTLA4 (MIM 123890), and ICOS (MIM 604558). May participate in the regulation of T-cell-mediated immune response. May play a protective role in tumor cells by inhibiting natural-killer mediated cell lysis as well as a role of marker for detection of neuroblastoma cells. May be involved in the development of acute and chronic transplant rejection and in the regulation of lymphocytic activity at mucosal surfaces. Could also play a key role in providing the placenta and fetus with a suitable immunological environment throughout pregnancy. Both isoform 1 and isoform 2 appear to be redundant in their ability to modulate CD4 T-cell responses. Isoform 2 is shown to enhance the induction of cytotoxic T-cells and selectively stimulates interferon gamma production in the presence of T-cell receptor signaling.

Applications Western Bloting: 1/500 - 1/2000.Immunofluorescence: 1/200 - 1/1000.Flow cytometry: 1/200 -1/400.ELISA: Propose dilution 1/10000.Not yet tested in other applications.Determining optimal working dilutions by titration test.

Limitations

- 1. Conjugates with brighter fluorochromes, like PE and APC, will have a greater separation than those with dyes like FITC and CyQ. When populations overlap, the percentage of positive cells using a selected marker can be affected by the choice of fluorescent label.
- 2. Use of monoclonal antibodies in patient treatment can interfere with antigen target recognition by this reagent. This should be taken into account when samples are analyzed from patients treated in this fashion. IQ Products has not characterized the effect of the presence of therapeutic antibodies on the performance of this reagent.
- 3. Reagents can be used in different combinations, therefore laboratories need to become familiar performance characteristics of each antibody in relation with the combined markers in normal and abnormal samples.
- 4. Reagent data performance is based on EDTA-treated blood. Reagent performance can be affected by the use of other anticoagulants.

IQP-1469 - CD276 Version 1

△ & / * □

Handling and Storage

Antibodies are supplied in 0.01 M sodium phosphate, 0.15 M NaCl; pH 7.3, 0.03% sodiumazide (NaN3) or as ascetic fluid containing 0.03% sodiumazide. Store the vials at 2-8 °C for a maximum of 2 weeks and store at -20°C for longer term storage. Monoclonal antibodies should be protected from prolonged exposure to light when conjugated with fluorochromes. Reagents are stable for the period shown on the vial label when stored properly.

Warranty

Products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied, which extend beyond the description on the label of the product. IQ Products is not liable for property damage, personal injury, or economic loss caused by the product.

Characterization

To ensure consistently high-quality reagents, each batch of monoclonal antibody is tested for conformance with characteristics of a standard reagent. Representative flow cytometric data is included in this data sheet.

Warning

All products contain sodiumazide. This chemical is poisonous and hazardous. Handling should be done by trained staff only.

References 1. Genome Res. 2004 Oct;14(10B):2121-7.

2. Cell Mol Immunol. 2005 Aug;2(4):307-11.

Explanation of used symbols

Consult instructions for use REF Catalogue number Sufficient for Δ Caution, consult accompanying document 米 Keep away from (sun)light Biological risks Temperature limitation (°C) RUO For Research Use Only LOT Batch code Use by yyyy-mm-dd

Manufacturer

bright fluorescence

EC REP

Authorized Representative in the European Community Conformité Européenne (European Conformity)

P F R C A	PURE FITC R-PE CyQ APC PerCP	Label - tandem purified material FITC PE PE-Cy5.18	Ex -max (nm) - 488 488, 532 488, 532 595, 633, 635, 647 488, 532	Em -max (nm) - 519 578 667 660 678
PCC	PerCP-Cy5.5		488, 532	695



IQ Products BV

Rozenburglaan 13a 9727 DL Groningen, The Netherlands

#31 (0)50 57 57 000 +31 (0)50 57 57 002

↑ Technical marketing@igproducts.nl
↑ Orders orders@igproducts.nl

www.igproducts.nl

IQP-1469 - CD276 Version 1