

PRODUCT INFORMATION SHEET

Mouse Monoclonal Antibody to STK11

Format: PURE RUO REF IQP-1507P Volume: 0.1ml

Description Ti

Clone 4H12

Mouse IgG1 **Isotype Specificity** STK11

Alternative names

PJS; LKB1; STK11

Species Human, Mouse & Monkey

Purified recombinant fragment of human STK11 expressed in E. Coli.

Immunogen

Mw 54kDa

Ascitic fluid containing 0.03% sodium azide.

Format

Summary

Essential role in G1 cell cycle arrest. Phosphorylates and activates members of the AMPK-related subfamily of protein kinases. Tumor suppressor.serine/threonine kinase 11, with two alternatively spliced isoforms, expressed in all tissues, strongly homolog of Xenopus early embryonic kinase 1 (XEEK1),tumor suppressor gene in hamartomas syndrome and in left sided colon carcinogenesis, mutated in sporadic testicular cancer, malignant melanomas and laryngeal tumors, playing a minor role in the development of ovarian carcinoma. LKB1 is a potential target for atherosclerosis and cancer and is shown to be mutated in patients with Peutz-Jeghers cancer syndrome. Tissue specificity: Ubiquitously expressed. Strongest expression in testis and fetal liver.

Applications Western Bloting: 1/500 - 1/2000.Flow cytometry: 1/200 - 1/400.ELISA: Propose dilution 1/10000.Not yet tested in other applications. Determining optimal working dilutions by titration

Limitations

- 1. Conjugates with brighter fluorochromes, like PE and APC, will have a greater separation than those with dyes like FITC and CyQ. When populations overlap, the percentage of positive cells using a selected marker can be affected by the choice of fluorescent label.
- 2. Use of monoclonal antibodies in patient treatment can interfere with antigen target recognition by this reagent. This should be taken into account when samples are analyzed from patients treated in this fashion. IQ Products has not characterized the effect of the presence of therapeutic antibodies on the performance of this reagent.
- 3. Reagents can be used in different combinations, therefore laboratories need to become familiar performance characteristics of each antibody in relation with the combined markers in normal and abnormal samples.
- 4. Reagent data performance is based on EDTA-treated blood. Reagent performance can be affected by the use of other anticoagulants.

IQP-1507 - STK11 Version 1

▲ ♦ 4 8

Handling and Storage

Antibodies are supplied in 0.01 M sodium phosphate, 0.15 M NaCl; pH 7.3, 0.03% sodiumazide (NaN3) or as ascetic fluid containing 0.03% sodiumazide. Store the vials at 2-8 °C for a maximum of 2 weeks and store at -20°C for longer term storage. Monoclonal antibodies should be protected from prolonged exposure to light when conjugated with fluorochromes. Reagents are stable for the period shown on the vial label when stored properly.

Warranty

Products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied, which extend beyond the description on the label of the product. IQ Products is not liable for property damage, personal injury, or economic loss caused by the product.

Characterization

To ensure consistently high-quality reagents, each batch of monoclonal antibody is tested for conformance with characteristics of a standard reagent. Representative flow cytometric data is included in this data sheet.

Warning

All products contain sodiumazide. This chemical is poisonous and hazardous. Handling should be done by trained staff only.

References 1. Sci Signal. 2009 Jul 21;2(80):ra35.

2. Sci Signal. 2009 Sep 1;2(86):pe55.

Explanation of used symbols

Consult instructions for use REF Catalogue number

Sufficient for

Δ Caution, consult accompanying document *

Keep away from (sun)light

8 Biological risks

Temperature limitation (°C)

RUO For Research Use Only

Batch code

Use by yyyy-mm-dd

Manufacturer Authorized Representative in the European Communit

Conformité Européenne (European Conformity)

P F R C A	PURE FITC R-PE CyQ APC	Label - tandem purified material FITC PE PE-Cy5.18	Ex -max (nm) - 488 488, 532 488, 532 595, 633, 635, 647	Em -max (nm) - 519 578 667 660
PC	PerCP		488, 532	678
PCC	PerCP-Cy5.5		488, 532	695



LOT

EC REP

IQ Products BV

Rozenburglaan 13a 9727 DL Groningen, The Netherlands

+31 (0)50 57 57 000 +31 (0)50 57 57 002

marketing@igproducts.nl Technical

Orders orders@igproducts.nl

www.igproducts.nl

IQP-1507 - STK11 Version 1