

PRODUCT INFORMATION SHEET

Mouse Monoclonal Antibody to CTCF

Format: PURE RUO REF IQP-1516P Volume: 0.1ml

Description \prod i

Clone 1D11

Mouse IgG1 **Isotype**

Specificity CTCF

Alternative names

CTCF

Species Human, Monkey

Immunogen Purified recombinant fragment of human CTCF expressed in E. Coli.

Mw 83kDa

Format Ascitic fluid containing 0.03% sodium azide.

Summary

This gene is a member of the BORIS + CTCF gene family and encodes a transcriptional regulator protein with 11 highly conserved zinc finger (ZF) domains. This nuclear protein is able to use different combinations of the ZF domains to bind different DNA target sequences and proteins. Depending upon the context of the site, the protein can bind a histone acetyltransferase (HAT)containing complex and function as a transcriptional activator or bind a histone deacetylase (HDAC)-containing complex and function as a transcriptional repressor. If the protein is bound to a transcriptional insulator element, it can block communication between enhancers and upstream promoters, thereby regulating imprinted expression. Mutations in this gene have been associated with invasive breast cancers, prostate cancers, and Wilms' tumors. Tissue specificity: Ubiquitous. Absent in primary spermatocytes.

Applications Western Bloting: 1/500 - 1/2000.ELISA: Propose dilution 1/10000.Not yet tested in other applications. Determining optimal working dilutions by titration test.

Limitations

- 1. Conjugates with brighter fluorochromes, like PE and APC, will have a greater separation than those with dyes like FITC and CyQ. When populations overlap, the percentage of positive cells using a selected marker can be affected by the choice of fluorescent label.
- 2. Use of monoclonal antibodies in patient treatment can interfere with antigen target recognition by this reagent. This should be taken into account when samples are analyzed from patients treated in this fashion. IO Products has not characterized the effect of the presence of therapeutic antibodies on the performance of this reagent.
- 3. Reagents can be used in different combinations, therefore laboratories need to become familiar performance characteristics of each antibody in relation with the combined markers in normal and abnormal samples.
- 4. Reagent data performance is based on EDTA-treated blood. Reagent performance can be affected by the use of other anticoagulants.

IOP-1516 - CTCF Version 1

△ ⊗ / * □

Handling and Storage

Antibodies are supplied in 0.01 M sodium phosphate, 0.15 M NaCl; pH 7.3, 0.03% sodiumazide (NaN3) or as ascetic fluid containing 0.03% sodiumazide. Store the vials at 2-8 °C for a maximum of 2 weeks and store at -20°C for longer term storage. Monoclonal antibodies should be protected from prolonged exposure to light when conjugated with fluorochromes. Reagents are stable for the period shown on the vial label when stored properly.

Warranty

Products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied, which extend beyond the description on the label of the product. IQ Products is not liable for property damage, personal injury, or economic loss caused by the product.

Characterization

To ensure consistently high-quality reagents, each batch of monoclonal antibody is tested for conformance with characteristics of a standard reagent. Representative flow cytometric data is included in this data sheet.

Warning

All products contain sodiumazide. This chemical is poisonous and hazardous. Handling should be done by trained staff only.

References 1. J Mol Med. 2008 Sep;86(9):1057-66.

- 2. J Exp Med. 2008 Apr 14;205(4):785-98.
- 3. EMBO J. 2008 Feb 20;27(4):654-66.

Explanation of used symbols



Consult instructions for use

REF

Catalogue number

Sufficient for

Caution, consult accompanying document

Keep away from (sun)light

Biological risks

Temperature limitation (°C)

RUO

For Research Use Only



Batch code
Use by yyy

Use by yyyy-mm-dd Manufacturer bright fluorescence



Authorized Representative in the European Community Conformité Européenne (European Conformity)

		Label - tandem	Ex -max (nm)	Em -max (nm)
Р	PURE	purified material	=	-
F	FITC	FITC	488	519
R	R-PE	PE	488, 532	578
С	CyQ	PE-Cy5.18	488, 532	667
Α	APC	•	595, 633, 635, 647	660
PC	PerCP		488, 532	678
PCC	PerCP-Cy5.5		488, 532	695
	,		,	



IQ Products BV

Rozenburglaan 13a 9727 DL Groningen, The Netherlands

** +31 (0)50 57 57 000 ** +31 (0)50 57 57 002

⁴ Technical <u>marketing@iqproducts.nl</u>⁴ Orders <u>orders@iqproducts.nl</u>

<u>www.iqproducts.nl</u>

IQP-1516 - CTCF Version 1