

PRODUCT INFORMATION SHEET

Mouse Monoclonal Antibody to TNNI2

Format: PURE RUO REF IQP-1646P Volume: 0.1ml

Description

Clone 2F12A8

Isotype Mouse IgG1

Specificity TNNI2

Alternative names

DA2B; FSSV; fsTnI; AMCD2B

Species Human

5 (6)

Immunogen

Purified recombinant fragment of human TNNI2 expressed in E. Coli.

Mw 21kDa

Format Ascitic fluid containing 0.03% sodium azide.

Summary

This gene encodes a fast-twitch skeletal muscle protein, a member of the troponin I gene family, and a component of the troponin complex including troponin T, troponin C and troponin I subunits. The troponin complex, along with tropomyosin, is responsible for the calcium-dependent regulation of striated muscle contraction. Mouse studies show that this component is also present in vascular smooth muscle and may play a role in regulation of smooth muscle function. In addition to muscle tissues, this protein is found in corneal epithelium, cartilage where it is an inhibitor of angiogenesis to inhibit tumor growth and metastasis, and mammary gland where it functions as a co-activator of estrogen receptor-related receptor alpha. This protein also suppresses tumor growth in human ovarian carcinoma. Mutations in this gene cause myopathy and distal arthrogryposis type 2B. Alternatively spliced transcript variants have been found for this gene.

Applications

Western Bloting: 1/500 - 1/2000.Immunohistochemistry: 1/200 - 1/1000.Flow cytometry: 1/200 - 1/400.ELISA: Propose dilution 1/10000.Not yet tested in other applications.Determining optimal working dilutions by titration test.

Limitations

- 1. Conjugates with brighter fluorochromes, like PE and APC, will have a greater separation than those with dyes like FITC and CyQ. When populations overlap, the percentage of positive cells using a selected marker can be affected by the choice of fluorescent label.
- 2. Use of monoclonal antibodies in patient treatment can interfere with antigen target recognition by this reagent. This should be taken into account when samples are analyzed from patients treated in this fashion. IQ Products has not characterized the effect of the presence of therapeutic antibodies on the performance of this reagent.
- 3. Reagents can be used in different combinations, therefore laboratories need to become familiar performance characteristics of each antibody in relation with the combined markers in normal and abnormal samples.
- Reagent data performance is based on EDTA-treated blood. Reagent performance can be affected by the use of other anticoagulants.

IOP-1646 - TNNI2 Version 1

▲ ♦ 4 8

Handling and Storage

Antibodies are supplied in 0.01 M sodium phosphate, 0.15 M NaCl; pH 7.3, 0.03% sodiumazide (NaN3) or as ascetic fluid containing 0.03% sodiumazide. Store the vials at 2-8 °C for a maximum of 2 weeks and store at -20°C for longer term storage. Monoclonal antibodies should be protected from prolonged exposure to light when conjugated with fluorochromes. Reagents are stable for the period shown on the vial label when stored properly.

Warranty

Products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied, which extend beyond the description on the label of the product. IQ Products is not liable for property damage, personal injury, or economic loss caused by the product.

Characterization

To ensure consistently high-quality reagents, each batch of monoclonal antibody is tested for conformance with characteristics of a standard reagent. Representative flow cytometric data is included in this data sheet.

Warning

RUO

LOT

All products contain sodiumazide. This chemical is poisonous and hazardous. Handling should be done by trained staff only.

ducts

t fluorescence

1. Am J Hum Genet. 2009 Nov;85(5):628-42. References 2. Cell Motil Cytoskeleton. 2008 Aug;65(8):652-61.

Explanation of used symbols



Ti Consult instructions for use REF Catalogue number Sufficient for

 \triangle Caution, consult accompanying document * Keep away from (sun)light 8

Biological risks

Temperature limitation (°C) For Research Use Only

Batch code

Use by yyyy-mm-dd Manufacturer

ECREP Authorized Representative in the European Community Conformité Européenne (European Conformity)

| P F | PURE FITC | Label - tandem purified material FITC | Ex -max (nm) - 488 | Em -max (nm) - 519 |
|--------|--------------|----------------------------------------------|--------------------------|---------------------------------|
| R | R-PE | PE | 488, 532 | 578 |
| С | CyQ | PE-Cy5.18 | 488, 532 | 667 |
| Α | APC | | 595, 633, 635, 647 | 660 |
| PC | PerCP | | 488, 532 | 678 |
| PCC | PerCP-Cy5.5 | | 488, 532 | 695 |

IQ Products BV

Rozenburglaan 13a 9727 DL Groningen, The Netherlands

+31 (0)50 57 57 000 +31 (0)50 57 57 002

Technical marketing@igproducts.nl Orders orders@igproducts.nl

www.igproducts.nl

IOP-1646 - TNNI2 Version 1