

PRODUCT INFORMATION SHEET

Mouse Monoclonal Antibody to MMP3

Format: PURE RUO REF IQP-1677P Volume: 0.1ml

Description

Clone 4F10

Isotype Mouse IgG1

MMP3

Specificity
Alternative

names SL-1; STMY; STR1; CHDS6; MMP-3; STMY1

Species Human

Immunogen Purified recombinant fragment of human MMP3 expressed in E. Coli.

Mw 54kDa

Format Purified antibody in PBS with 0.05% sodium azide and 0.5% protein stabilizer

Summary Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of

extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. This gene encodes an enzyme which degrades fibronectin, laminin, collagens III, IV, IX, and X, and cartilage proteoglycans. The enzyme is thought to be involved in wound repair, progression of atherosclerosis, and tumor initiation. The gene is part of a cluster of

MMP genes which localize to chromosome 11q22.3.

Applications Western Bloting: 1/500 - 1/2000.Flow cytometry: 1/200 - 1/400.ELISA: Propose dilution

1/10000.Not yet tested in other applications. Determining optimal working dilutions by titration

test.

Limitations

- 1. Conjugates with brighter fluorochromes, like PE and APC, will have a greater separation than those with dyes like FITC and CyQ. When populations overlap, the percentage of positive cells using a selected marker can be affected by the choice of fluorescent label.
- Use of monoclonal antibodies in patient treatment can interfere with antigen target recognition by this reagent. This should be taken into account when samples are analyzed from patients treated in this fashion. IQ Products has not characterized the effect of the presence of therapeutic antibodies on the performance of this reagent.
- 3. Reagents can be used in different combinations, therefore laboratories need to become familiar performance characteristics of each antibody in relation with the combined markers in normal and abnormal samples.
- 4. Reagent data performance is based on EDTA-treated blood. Reagent performance can be affected by the use of other anticoagulants.

IQP-1677 – MMP3 Version 1

△ ♦ / * □

Handling and Storage

Antibodies are supplied in 0.01 M sodium phosphate, 0.15 M NaCl; pH 7.3, 0.03% sodiumazide (NaN3) or as ascetic fluid containing 0.03% sodiumazide. Store the vials at 2-8 °C for a maximum of 2 weeks and store at -20°C for longer term storage. Monoclonal antibodies should be protected from prolonged exposure to light when conjugated with fluorochromes. Reagents are stable for the period shown on the vial label when stored properly.

Warranty

Products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied, which extend beyond the description on the label of the product. IQ Products is not liable for property damage, personal injury, or economic loss caused by the product.

Characterization

To ensure consistently high-quality reagents, each batch of monoclonal antibody is tested for conformance with characteristics of a standard reagent. Representative flow cytometric data is included in this data sheet.

Warning

RUO

EC REP

All products contain sodiumazide. This chemical is poisonous and hazardous. Handling should be done by trained staff only.

References 1. Int J Cancer. 2009 May 1;124(9):2172-8. 2. Cancer Epidemiol Biomarkers Prev. 2008 Dec;17(12):3551-7.

Explanation of used symbols

Consult instructions for use

REF

Sufficient for

Caution, consult accompanying document

Keep away from (sun)light

Biological risks
Temperature limitation (°C)

For Research Use Only

LOT Batch code

Use by yyyy-mm-dd Manufacturer

Authorized Representative in the European Community Conformité Européenne (European Conformity)

| P F R C A | PURE FITC R-PE CyQ APC PerCP | Label - tandem purified material FITC PE PE-Cy5.18 | Ex -max (nm) - 488 488, 532 488, 532 595, 633, 635, 647 488, 532 | Em -max (nm) - 519 578 667 660 678 |
|-----------------------|---|---|--|------------------------------------|
| PCC | PerCP-Cy5.5 | | 488, 532 | 695 |



IQ Products BV

Rozenburglaan 13a 9727 DL Groningen, The Netherlands

** +31 (0)50 57 57 000 ** +31 (0)50 57 57 002

✓ Technical marketing@iqproducts.nl✓ Orders orders@iqproducts.nl

<u>www.iqproducts.nl</u>

IQP-1677 – MMP3 Version 1