

## **PRODUCT INFORMATION SHEET**

|                          | Format: PURE   | RUO  | REF   | IQP-1678P   | Volume:  | 0.1ml                              |  |
|--------------------------|--|--|---|---|--|------------------------------------|--|
|                          |  |  |   |   |  |                                    |  |
| Ĩ                        | Description  |  |   |   |  |                                    |  |
| Clone                    | 6G5  |  |   |   |  |                                    |  |
| sotype                   | Mouse IgG1   |  |   |   |  |                                    |  |
| pecificity               | CYP1A1   |  |   |   |  |                                    |  |
| lternative<br>ames       | AHH; AHRR; CP11; CYP1; P1-450; P450-C; P450DX  |  |   |   |  |                                    |  |
| pecies                   | Human  |  |   |   |  |                                    |  |
| mmunogen                 | Purified recombinant fragment of human CYP1A1 expressed in E. Coli.  |  |   |   |  |                                    |  |
| 1w<br><sup>:</sup> ormat | 58kDa<br>Purified antibody in PBS with 0.05% sodium azide and 0.5% protein stabilizer  |  |   |   |  |                                    |  |
| ummary                   | This gene, CYP1A1, encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and its expression is induced by some polycyclic aromatic hydrocarbons (PAHs), some of which are found in cigarette smoke. The enzyme's endogenous substrate is unknown; however, it is able to metabolize some PAHs to carcinogenic intermediates. The gene has been associated with lung cancer risk. A related family member, CYP1A2, is located approximately 25 kb away from CYP1A1 on chromosome 15. |  |   |   |  |                                    |  |
| pplications              | Western Bloting: 1/50<br>1/10000.Not yet teste<br>test.  |  |   |   |  |                                    |  |
| imitations               | <ol> <li>Conjugates with t<br/>those with dyes li</li> </ol>   |  |   |   |  |                                    |  |
|                          | using a selected r<br>2. Use of monoclona<br>by this reagent. T<br>treated in this fas   | narker can be af<br>Il antibodies in pa<br>his should be tal<br>hion. IQ Product | fected by th<br>atient treatr<br>ken into acc<br>s has not ch | e choice of fluore<br>ment can interfere<br>ount when sampl<br>naracterized the e | scent label.<br>e with antigen<br>es are analyze | target recognit<br>d from patients |  |
|                          | therapeutic antibo<br>3. Reagents can be<br>familiar performa  | used in different<br>nce characteristic  | combinatio  | ns, therefore labo  |  |                                    |  |
|                          | normal and abnor   | mal samples.   |   |   |  |                                    |  |

# ⚠ 💩 🔏 🛣 🗳

## Handling and Storage

Antibodies are supplied in 0.01 M sodium phosphate, 0.15 M NaCl; pH 7.3, 0.03% sodiumazide (NaN3) or as ascetic fluid containing 0.03% sodiumazide. Store the vials at 2-8 °C for a maximum of 2 weeks and store at -20°C for longer term storage. Monoclonal antibodies should be protected from prolonged exposure to light when conjugated with fluorochromes. Reagents are stable for the period shown on the vial label when stored properly.

**Warranty** Products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied, which extend beyond the description on the label of the product. IQ Products is not liable for property damage, personal injury, or economic loss caused by the product.

#### Characterization

- To ensure consistently high-quality reagents, each batch of monoclonal antibody is tested for conformance with characteristics of a standard reagent. Representative flow cytometric data is included in this data sheet.
- **Warning** All products contain sodiumazide. This chemical is poisonous and hazardous. Handling should be done by trained staff only.

**References** 1. Cancer Res. 2009 Apr 1;69(7):2956-65. 2. Drug Metab Lett. 2009 Jan;3(1):18-27.

### Explanation of used symbols

| (li)       | Consult instructions for use                        |
|------------|---|
| 「追<br>REF  | Catalogue number                                    |
| ¥          | Sufficient for                                      |
|            | Caution, consult accompanying document              |
| *          | Keep away from (sun)light                           |
| & /        | Biological risks                                    |
| X          | Temperature limitation (°C)                         |
| RUO        | For Research Use Only                               |
| RUO<br>LOT | Batch code  |
| 2          | Use by yyyy-mm-dd                                   |
|            | Manufacturer OFICIAL TIUORESCENCE                   |
| EC REP     | Authorized Representative in the European Community |
| CE         | Conformité Européenne (European Conformity)         |
|            |   |

| P<br>F<br>C<br>A | PURE<br>FITC<br>R-PE<br>CyQ<br>APC<br>PerCP | Label - tandem<br>purified material<br>FITC<br>PE<br>PE-Cy5.18 | <b>Ex -max (nm)</b><br>-<br>488<br>488, 532<br>488, 532<br>595, 633, 635, 647<br>488, 532 | <b>Em -max (nm)</b><br>-<br>519<br>578<br>667<br>660<br>678 |
|------------------|---|--|---|---|
| PC               | PerCP                                       |  | 488, 532  | 678   |
| PCC              | PerCP-Cy5.5                                 |  | 488, 532  | 695   |

IQ Products BV

Rozenburglaan 13a 9727 DL Groningen, The Netherlands

- +31 (0)50 57 57 000
- ∃ +31 (0)50 57 57 002
- Technical <u>marketing@iqproducts.nl</u>
- Orders <u>orders@iqproducts.nl</u>
- <u>www.iqproducts.nl</u>