

PRODUCT INFORMATION SHEET

Mouse Monoclonal Antibody to E2F1

Format: PURE RUO REF IQP-1752P Volume: 0.1ml

Description

Clone 8G9

Isotype Mouse IgG1

Specificity E2F1

Alternative names

RBP3; E2F-1; RBAP1; RBBP3

Species Human, Mouse

Purified recombinant fragment of human E2F1 (AA: 69-223) expressed in E. Coli.

Immunogen

Mw 46.9kDa

Format Purified antibody in PBS with 0.05% sodium azide and 0.5% protein stabilizer

Summary

The protein encoded by this gene is a member of the E2F family of transcription factors. The E2F family plays a crucial role in the control of cell cycle and action of tumor suppressor proteins and is also a target of the transforming proteins of small DNA tumor viruses. The E2F proteins contain several evolutionally conserved domains found in most members of the family. These domains include a DNA binding domain, a dimerization domain which determines interaction with the differentiation regulated transcription factor proteins (DP), a transactivation domain enriched in acidic amino acids, and a tumor suppressor protein association domain which is embedded within the transactivation domain. This protein and another 2 members, E2F2 and E2F3, have an additional cyclin binding domain. This protein binds preferentially to retinoblastoma protein pRB in a cell-cycle dependent manner. It can mediate both cell proliferation and p53-dependent/independent apoptosis.

Applications

Limitations

- 1. Conjugates with brighter fluorochromes, like PE and APC, will have a greater separation than those with dyes like FITC and CyQ. When populations overlap, the percentage of positive cells using a selected marker can be affected by the choice of fluorescent label.
- Use of monoclonal antibodies in patient treatment can interfere with antigen target recognition by this reagent. This should be taken into account when samples are analyzed from patients treated in this fashion. IQ Products has not characterized the effect of the presence of therapeutic antibodies on the performance of this reagent.
- 3. Reagents can be used in different combinations, therefore laboratories need to become familiar performance characteristics of each antibody in relation with the combined markers in normal and abnormal samples.
- 4. Reagent data performance is based on EDTA-treated blood. Reagent performance can be affected by the use of other anticoagulants.

IOP-1752 - E2F1 Version 1

▲ ♦ / * 2

Handling and Storage

Antibodies are supplied in 0.01 M sodium phosphate, 0.15 M NaCl; pH 7.3, 0.03% sodiumazide (NaN3) or as ascetic fluid containing 0.03% sodiumazide. Store the vials at 2-8 °C for a maximum of 2 weeks and store at -20°C for longer term storage. Monoclonal antibodies should be protected from prolonged exposure to light when conjugated with fluorochromes. Reagents are stable for the period shown on the vial label when stored properly.

Warranty

Products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied, which extend beyond the description on the label of the product. IQ Products is not liable for property damage, personal injury, or economic loss caused by the product.

Characterization

To ensure consistently high-quality reagents, each batch of monoclonal antibody is tested for conformance with characteristics of a standard reagent. Representative flow cytometric data is included in this data sheet.

Warning

All products contain sodiumazide. This chemical is poisonous and hazardous. Handling should be done by trained staff only.

ducts

1.Cancer Res. 2010 Dec 1;70(23):9711-20. 2.Mol Cancer Ther. 2010 May;9(5):1265-73.

Explanation of used symbols



Ţi Consult instructions for use REF Catalogue number Sufficient for Δ Caution, consult accompanying document * Keep away from (sun)light Biological risks

Temperature limitation (°C) RUO For Research Use Only

LOT Batch code

EC REP

Use by yyyy-mm-dd Manufacturer

fluorescence Authorized Representative in the European Community Conformité Européenne (European Conformity)

P F R C A	PURE FITC R-PE CyQ APC PerCP	Label - tandem purified material FITC PE PE-Cy5.18	Ex -max (nm) - 488 488, 532 488, 532 595, 633, 635, 647 488, 532	Em -max (nm) - 519 578 667 660 678
PCC	PerCP-Cy5.5		488, 532	695



IQ Products BV

Rozenburglaan 13a 9727 DL Groningen, The Netherlands

+31 (0)50 57 57 000 +31 (0)50 57 57 002

Technical <u>marketing@iqproducts.nl</u> Orders orders@igproducts.nl

www.igproducts.nl

IQP-1752 - E2F1 Version 1