

PRODUCT INFORMATION SHEET

Mouse Monoclonal Antibody to CSNK2A2							
	Format: PURE	RUO	REF	IQP-1773P	Volume:	0.1ml	
	Description						
Clone	1E1						
Isotype	Mouse IgG1						
Specificity	CSNK2A2						
Alternative names	CK2A2; CSNK2A1; FLJ43934						
Species	Human, Mouse						
Immunogen	Purified recombinant fragment of human CSNK2A2 (AA: 194-350) expressed in E. Coli.						
Mw	41.2kDa						
Format	Ascitic fluid containing 0.03% sodium azide.						
Summary	Casein kinase II (CK2) is a constitutively active, ubiquitously expressed serine/threonine protein kinasethat is thought to have a regulatory function in cell proliferation, cell differentiation and apoptosis. CK2functions as a tetrameric complex consisting of two regulatory beta-subunits and two catalytic units (alphaand alpha') in a homomeric or heteromeric conformation. Whilst the alpha- and alpha'-subunits arecatalytically identical, proteins that regulate CK2, such as cdc2 and Hsp90, preferentially bind to thealpha and not the alpha'-subunit. CK2 can phosphorylate a number of key intracellular signaling proteinsimplicated in tumor suppression (p53 and PTEN) and tumorigenesis (myc, jun, NF-kappaB). CK2 is also thoughtto influence Wnt signaling via beta-catenin phosphorylation and the PI 3-K signaling pathway via thephosphorylation of Akt.						
Applications						4 c	
Limitations	 Conjugates with brighter fluorochromes, like PE and APC, will have a greater separation t those with dyes like FITC and CyQ. When populations overlap, the percentage of positive using a selected marker can be affected by the choice of fluorescent label. Use of monoclonal antibodies in patient treatment can interfere with antigen target recog by this reagent. This should be taken into account when samples are analyzed from patie treated in this fashion. IQ Products has not characterized the effect of the presence of therapeutic antibodies on the performance of this reagent. 						

- 3. Reagents can be used in different combinations, therefore laboratories need to become familiar performance characteristics of each antibody in relation with the combined markers in normal and abnormal samples.
- normal and abnormal samples.Reagent data performance is based on EDTA-treated blood. Reagent performance can be affected by the use of other anticoagulants.

⚠ 💩 🔏 🛣 🎴

Handling and Storage

Antibodies are supplied in 0.01 M sodium phosphate, 0.15 M NaCl; pH 7.3, 0.03% sodiumazide (NaN3) or as ascetic fluid containing 0.03% sodiumazide. Store the vials at 2-8 °C for a maximum of 2 weeks and store at -20°C for longer term storage. Monoclonal antibodies should be protected from prolonged exposure to light when conjugated with fluorochromes. Reagents are stable for the period shown on the vial label when stored properly.

Warranty Products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied, which extend beyond the description on the label of the product. IQ Products is not liable for property damage, personal injury, or economic loss caused by the product.

Characterization

- To ensure consistently high-quality reagents, each batch of monoclonal antibody is tested for conformance with characteristics of a standard reagent. Representative flow cytometric data is included in this data sheet.
- **Warning** All products contain sodiumazide. This chemical is poisonous and hazardous. Handling should be done by trained staff only.

References 1.Eur J Immunol. 2009 Jan;39(1):267-79. 2.Cell Mol Life Sci. 2009 Jan;66(2):339-49.

Explanation of used symbols

	Consult instructions for use
REF	Catalogue number
V	Sufficient for
	Caution, consult accompanying document
*	Keep away from (sun)light
l de	Biological risks
X /	Temperature limitation (°C)
RUO LOT	For Research Use Only
LOT	Batch code
	Use by yyyy-mm-dd
1	Manufacturer
EC REP	Authorized Representative in the European Community
CE	Conformité Européenne (European Conformity)

P F R	PURE FITC R-PE	Label - tandem purified material FITC PE	Ex -max (nm) - 488 488, 532	Em -max (nm) - 519 578
C	CyQ	PE-Cy5.18	488, 532	667
A	APC		595, 633, 635, 647	660
PC	PerCP		488, 532	678
PCC	PerCP-Cy5.5		488, 532	695

IQ Products BV

Rozenburglaan 13a 9727 DL Groningen, The Netherlands

+31 (0)50 57 57 000

- ₼ +31 (0)50 57 57 002
- Technical <u>marketing@iqproducts.nl</u>
- Orders <u>orders@iqproducts.nl</u>
- <u>www.iqproducts.nl</u>