

PRODUCT INFORMATION SHEET

Mouse Monoclonal Antibody to DKK3

Format: PURE RUO REF IQP-1803P Volume: 0.1mg

Description

Clone 8A5C6

Isotype Mouse IgG1

Specificity DKK3

Alternative names

RIG; REIC

Species Human,

Immunogen Purified recombinant fragment of human DKK3 (AA: 91-350) expressed in E. Coli.

Mw 38.3kDa

Format Purified antibody in PBS with 0.05% sodium azide and 0.5% protein stabilizer

Summary This gene encodes a protein that is a member of the dickkopf family. The secreted protein contains

two cysteine rich regions and is involved in embryonic development through its interactions with the Wnt signaling pathway. The expression of this gene is decreased in a variety of cancer cell lines and it may function as a tumor suppressor gene. Alternative splicing results in multiple

transcript variants encoding the same protein.

Applications ELISA: 1/10000; WB: 1/500 - 1/2000; IHC: 1/200 - 1/1000; FCM: 1/200 - 1/400

Limitations

- 1. Conjugates with brighter fluorochromes, like PE and APC, will have a greater separation than those with dyes like FITC and CyQ. When populations overlap, the percentage of positive cells using a selected marker can be affected by the choice of fluorescent label.
- 2. Use of monoclonal antibodies in patient treatment can interfere with antigen target recognition by this reagent. This should be taken into account when samples are analyzed from patients treated in this fashion. IQ Products has not characterized the effect of the presence of therapeutic antibodies on the performance of this reagent.
- 3. Reagents can be used in different combinations, therefore laboratories need to become familiar performance characteristics of each antibody in relation with the combined markers in normal and abnormal samples.
- 4. Reagent data performance is based on EDTA-treated blood. Reagent performance can be affected by the use of other anticoagulants.

IQP-1803 – DKK3 Version 1

△ ⊗ / * □

Handling and Storage

Antibodies are supplied in 0.01 M sodium phosphate, 0.15 M NaCl; pH 7.3, 0.03% sodiumazide (NaN3) or as ascetic fluid containing 0.03% sodiumazide. Store the vials at 2-8 °C for a maximum of 2 weeks and store at -20°C for longer term storage. Monoclonal antibodies should be protected from prolonged exposure to light when conjugated with fluorochromes. Reagents are stable for the period shown on the vial label when stored properly.

Warranty

Products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied, which extend beyond the description on the label of the product. IQ Products is not liable for property damage, personal injury, or economic loss caused by the product.

Characterization

To ensure consistently high-quality reagents, each batch of monoclonal antibody is tested for conformance with characteristics of a standard reagent. Representative flow cytometric data is included in this data sheet.

Warning

All products contain sodiumazide. This chemical is poisonous and hazardous. Handling should be done by trained staff only.

References 1.Exp Dermatol. 2011 Mar;20(3):273-7. 2.Thromb Haemost. 2011 Jan;105(1):72-80.

Explanation of used symbols

Consult instructions for use

REF

Catalogue number

Sufficient for

Caution, consult accompanying document

Keep away from (sun)light

Biological risks

Temperature limitation (°C)
For Research Use Only

LOT Batch code

Use by yyyy-mm-dd

Manufacturer

Authorized Representative in the European Community Conformité Européenne (European Conformity)

		Label - tandem	Ex -max (nm)	Em -max (nm)
Р	PURE	purified material	-	-
F	FITC	FITC	488	519
R	R-PE	PE	488, 532	578
С	CyQ	PE-Cy5.18	488, 532	667
Α	APC	•	595, 633, 635, 647	660
PC	PerCP		488, 532	678
PCC	PerCP-Cy5.5		488, 532	695



EC REP

IQ Products BV

Rozenburglaan 13a 9727 DL Groningen, The Netherlands

★ +31 (0)50 57 57 000★ +31 (0)50 57 57 002

Technical marketing@iqproducts.nlOrders orders@iqproducts.nl

www.iqproducts.nl

IQP-1803 – DKK3 Version 1