

## PRODUCT INFORMATION SHEET

### Mouse Monoclonal Antibody to SNCG (breast cancer-specific protein 1)

Format: PURE      RUO      REF      IQP-1073P      Volume: 0.1ml



#### Description

<b>Clone</b>	1H10D2
<b>Isotype</b>	Mouse IgG1
<b>Specificity</b>	SNCG (breast cancer-specific protein 1)
<b>Alternative names</b>	SR; BCSG1
<b>Species</b>	Human
<b>Immunogen</b>	Purified recombinant fragment of SNCG expressed in E. Coli.
<b>Mw</b>	
<b>Format</b>	Purified antibody in PBS containing 0.03% sodium azide.

**Summary** SNCG(also designated gamma-synuclein or breast cancer-specific protein 1),with 127-amino acid protein(about 14kDa), belongs to the synuclein family, which also includes alpha- and beta-synuclein.Three synucleins are located in the neuronal cytosol and enriched in presynaptic terminals,while SNCG is also expressed in many other non-neuronal tissues. SNCG is abnormally expressed in a high percentage of tumor tissues of diversified cancer types, including liver, esophagus, colon, gastric, lung, prostate, cervical, and breast cancer, but rarely expressed in tumor-matched nonneoplastic adjacent tissues. High levels of SNCG have been identified in advanced breast carcinomas suggesting a correlation between overexpression of SNCG and breast tumor development.

**Applications** Western Blotting: 1/500 - 1/2000.Immunohistochemistry: 1/200 - 1/1000.ELISA: Propose dilution 1/10000.Not yet tested in other applications.Determining optimal working dilutions by titration test.

#### Limitations

1. Conjugates with brighter fluorochromes, like PE and APC, will have a greater separation than those with dyes like FITC and CyQ. When populations overlap, the percentage of positive cells using a selected marker can be affected by the choice of fluorescent label.
2. Use of monoclonal antibodies in patient treatment can interfere with antigen target recognition by this reagent. This should be taken into account when samples are analyzed from patients treated in this fashion. IQ Products has not characterized the effect of the presence of therapeutic antibodies on the performance of this reagent.
3. Reagents can be used in different combinations, therefore laboratories need to become familiar performance characteristics of each antibody in relation with the combined markers in normal and abnormal samples.
4. Reagent data performance is based on EDTA-treated blood. Reagent performance can be affected by the use of other anticoagulants.



### Handling and Storage

Antibodies are supplied in 0.01 M sodium phosphate, 0.15 M NaCl; pH 7.3, 0.03% sodiumazide (NaN<sub>3</sub>) or as ascetic fluid containing 0.03% sodiumazide. Store the vials at 2-8 °C for a maximum of 2 weeks and store at -20°C for longer term storage. Monoclonal antibodies should be protected from prolonged exposure to light when conjugated with fluorochromes. Reagents are stable for the period shown on the vial label when stored properly.

### Warranty

Products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied, which extend beyond the description on the label of the product. IQ Products is not liable for property damage, personal injury, or economic loss caused by the product.

### Characterization

To ensure consistently high-quality reagents, each batch of monoclonal antibody is tested for conformance with characteristics of a standard reagent. Representative flow cytometric data is included in this data sheet.

### Warning

All products contain sodiumazide. This chemical is poisonous and hazardous. Handling should be done by trained staff only.

- 
- References** 1. Haiyan Liu, Wei Liu, Yinwei Wu. Cancer Res. 2005 Sep 1;65(17):7635-43.  
2. Irina Surgucheva , Belinda McMahon , Andrei Surguchov. Cell Motil Cytoskeleton. 2006 May 26.
- 

### Explanation of used symbols



REF



RUO

LOT



EC REP



Consult instructions for use

Catalogue number

Sufficient for

Caution, consult accompanying document

Keep away from (sun)light

Biological risks

Temperature limitation (°C)

For Research Use Only

Batch code

Use by yyyy-mm-dd

Manufacturer

Authorized Representative in the European Community

Conformité Européenne (European Conformity)



**IQ Products**  
bright fluorescence

		<b>Label - tandem</b>	<b>Ex -max (nm)</b>	<b>Em -max (nm)</b>
P	PURE	purified material	-	-
F	FITC	FITC	488	519
R	R-PE	PE	488, 532	578
C	CyQ	PE-Cy5.18	488, 532	667
A	APC		595, 633, 635, 647	660
PC	PerCP		488, 532	678
PCC	PerCP-Cy5.5		488, 532	695



IQ Products BV

Rozenburglaan 13a  
9727 DL Groningen, The Netherlands

+31 (0)50 57 57 000

+31 (0)50 57 57 002

Technical [marketing@iqproducts.nl](mailto:marketing@iqproducts.nl)

Orders [orders@iqproducts.nl](mailto:orders@iqproducts.nl)

[www.iqproducts.nl](http://www.iqproducts.nl)