

PRODUCT INFORMATION SHEET

Mouse Monoclonal Antibody to Oct4

Format: PURE RUO REF IQP-1246P Volume: 100µl

Description

Clone 1D10H6

Isotype Mouse IgG1

Specificity Oct4

Alternative names

OCT3; OTF3; OTF4; MGC22487; POU5F1

Species Human

Immunogen Purified recombinant fragment of Oct4 (aa193-360) expressed in E. Coli.

Mw 38.5kDa

Format Ascitic fluid containing 0.03% sodium azide.

Summary

Oct4: octamer-binding transcription factor-4 (Oct-4, Otf-4) and Oct-3/4, also known as POU5F1(POU class 5 homeobox 1), octamerbinding transcription factor-3 (Oct-3, Otf-3), modulates embryonic stem (ES) cell populations by influencing lineage commitment. Entrez Protein NP_002692. Oct-3/4 sustains stem-cell selfrenewal and differentiation pathways. Transcription factors containing the POU homeodomain regulate tissue-specific gene expression in lymphoid and pituitary differentiation and in early mammalian development. Oct-3/4 is capable of inducing rapid proliferation and tumorigenic properties of ES cells through activation of the UTF1 gene. In humans, two Oct-3/4 isoforms contribute to influencing the undifferentiated phenotype of ES cells. Oct-3/4 pseudogenes localizing to human chromosomes 10 and 8 are reported to be transcribed in certain cancer cell lines and tissues.

Applications ELISA: 1/10000; WB: 1/500 - 1/2000

Limitations

- 1. Conjugates with brighter fluorochromes, like PE and APC, will have a greater separation than those with dyes like FITC and CyQ. When populations overlap, the percentage of positive cells using a selected marker can be affected by the choice of fluorescent label.
- Use of monoclonal antibodies in patient treatment can interfere with antigen target recognition by this reagent. This should be taken into account when samples are analyzed from patients treated in this fashion. IQ Products has not characterized the effect of the presence of therapeutic antibodies on the performance of this reagent.
- 3. Reagents can be used in different combinations, therefore laboratories need to become familiar performance characteristics of each antibody in relation with the combined markers in normal and abnormal samples.
- 4. Reagent data performance is based on EDTA-treated blood. Reagent performance can be affected by the use of other anticoagulants.

△ ⊗ / * 2

Handling and Storage

Store the vials at 2-8 °C for a maximum of 2 weeks and store at -20°C for longer term storage. Monoclonal antibodies should be protected from prolonged exposure to light when conjugated with fluorochromes. Reagents are stable for the period shown on the vial label when stored properly.

IQP-1246 - Oct4 Version 2

Warranty

Products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied, which extend beyond the description on the label of the product. IQ Products is not liable for property damage, personal injury, or economic loss caused by the product.

Characterization

To ensure consistently high-quality reagents, each batch of monoclonal antibody is tested for conformance with characteristics of a standard reagent.

Warning

All products contain sodiumazide. This chemical is poisonous and hazardous. Handling should be done by trained staff only.

References

- 1. Cells Tissues Organs. 2006;184(3-4):105-16.
- 2. Int J Cancer. 2007 Apr 1;120(7):1598-602.

Explanation of used symbols



		Label - tandem	Ex -max (nm)	Em -max (nm)
P	PURE	purified material	-	-
F	FITC	FITC	488	519
R	R-PE	PE PE	488, 532	578
С	CyQ	PE-Cy5.18	488, 532	667
Α	APC	_	595, 633, 635, 647	660
PC	PerCP		488, 532	678
PCC	PerCP-Cy5.5		488, 532	695



IQ Products BV

Rozenburglaan 13a 9727 DL Groningen, The Netherlands

★ +31 (0)50 57 57 000★ +31 (0)50 57 57 002

Technical marketing@iqproducts.nlOrders orders@iqproducts.nl

<u>www.igproducts.nl</u>

IQP-1246 – Oct4 Version 2