

## PRODUCT INFORMATION SHEET

### Mouse Monoclonal Antibody to Pirh2

Format: PURE      RUO      REF      IQP-1338P      Volume: 0.1ml



#### Description

<b>Clone</b>	1H10
<b>Isotype</b>	Mouse IgG1
<b>Specificity</b>	Pirh2
<b>Alternative names</b>	ARNIP; CHIMP; RNF199; RCHY1
<b>Species</b>	Human, Rat
<b>Immunogen</b>	Purified recombinant fragment of human Pirh2 expressed in E. Coli.
<b>Mw</b>	30kDa; 60kDa (homodimer)
<b>Format</b>	Ascitic fluid containing 0.03% sodium azide.

**Summary** Pirh 2 (P53 induced RING-H2 protein), also known as RCHY1, it forms dimers through its N- and C-terminus in cells. The Pirh2 has ubiquitin-protein ligase activity and it binds with p53 and promotes the ubiquitin-mediated proteosomal degradation of p53. The Pirh2 is oncogenic because loss of p53 function contributes directly to malignant tumor development. Pirh2 expression decreases the level of p53, and a decrease of endogenous Pirh2 expression increases p53 levels. Pirh2 is therefore considered, together with MDM2, to act as a negative regulator of p53 function.

**Applications** Western Blotting: 1/500 - 1/2000. Immunohistochemistry: 1/200 - 1/1000. Immunofluorescence: 1/200 - 1/1000. Flow cytometry: 1/200 - 1/400. ELISA: Propose dilution 1/10000. Not yet tested in other applications. Determining optimal working dilutions by titration test.

#### Limitations

1. Conjugates with brighter fluorochromes, like PE and APC, will have a greater separation than those with dyes like FITC and CyQ. When populations overlap, the percentage of positive cells using a selected marker can be affected by the choice of fluorescent label.
2. Use of monoclonal antibodies in patient treatment can interfere with antigen target recognition by this reagent. This should be taken into account when samples are analyzed from patients treated in this fashion. IQ Products has not characterized the effect of the presence of therapeutic antibodies on the performance of this reagent.
3. Reagents can be used in different combinations, therefore laboratories need to become familiar performance characteristics of each antibody in relation with the combined markers in normal and abnormal samples.
4. Reagent data performance is based on EDTA-treated blood. Reagent performance can be affected by the use of other anticoagulants.



### Handling and Storage

Antibodies are supplied in 0.01 M sodium phosphate, 0.15 M NaCl; pH 7.3, 0.03% sodiumazide (NaN<sub>3</sub>) or as ascetic fluid containing 0.03% sodiumazide. Store the vials at 2-8 °C for a maximum of 2 weeks and store at -20°C for longer term storage. Monoclonal antibodies should be protected from prolonged exposure to light when conjugated with fluorochromes. Reagents are stable for the period shown on the vial label when stored properly.

### Warranty

Products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied, which extend beyond the description on the label of the product. IQ Products is not liable for property damage, personal injury, or economic loss caused by the product.

### Characterization

To ensure consistently high-quality reagents, each batch of monoclonal antibody is tested for conformance with characteristics of a standard reagent. Representative flow cytometric data is included in this data sheet.

### Warning

All products contain sodiumazide. This chemical is poisonous and hazardous. Handling should be done by trained staff only.

- 
- References**
1. Biochem Biophys Res Commun. 2007 Dec 14;364(2):344-50.
  2. J Natl Cancer Inst. 2004 Nov 17;96(22):1718-21.
  3. Exp Cell Res. 2006 Oct 15;312(17):3370-8.
  4. Cell. 2003 Mar 21;112(6):779-91.
- 

### Explanation of used symbols



Consult instructions for use



Catalogue number



Sufficient for



Caution, consult accompanying document



Keep away from (sun)light



Biological risks



Temperature limitation (°C)



For Research Use Only



Batch code



Use by yyyy-mm-dd



Manufacturer



Authorized Representative in the European Community



Conformité Européenne (European Conformity)

		<b>Label - tandem</b>	<b>Ex -max (nm)</b>	<b>Em -max (nm)</b>
P	PURE	purified material	-	-
F	FITC	FITC	488	519
R	R-PE	PE	488, 532	578
C	CyQ	PE-Cy5.18	488, 532	667
A	APC		595, 633, 635, 647	660
PC	PerCP		488, 532	678
PCC	PerCP-Cy5.5		488, 532	695

---



IQ Products BV  
Rozenburglaan 13a  
9727 DL Groningen, The Netherlands

- +31 (0)50 57 57 000
- +31 (0)50 57 57 002
- Technical [marketing@iqproducts.nl](mailto:marketing@iqproducts.nl)
- Orders [orders@iqproducts.nl](mailto:orders@iqproducts.nl)
- [www.iqproducts.nl](http://www.iqproducts.nl)