

## PRODUCT INFORMATION SHEET

### Mouse Monoclonal Antibody to UBE1L

Format: PURE      RUO      REF      IQP-1459P      Volume: 0.1ml

 Description	
<b>Clone</b>	5B10
<b>Isotype</b>	Mouse IgG1
<b>Specificity</b>	UBE1L
<b>Alternative names</b>	D8; UBE2; UBA1B; UBE1L; MGC12713; UBA7
<b>Species</b>	Human
<b>Immunogen</b>	Purified recombinant fragment of human UBE1L expressed in E. Coli.
<b>Mw</b>	112kDa
<b>Format</b>	Ascitic fluid containing 0.03% sodium azide.
<b>Summary</b>	<p>The modification of proteins with ubiquitin is an important cellular mechanism for targeting. Tissue specificity: Expressed in a variety of normal and tumor cell types, but is reduced in lung cancer cell lines abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes, or E1s, ubiquitin-conjugating enzymes, or E2s, and ubiquitin-protein ligases, or E3s. This gene encodes a member of the E1 ubiquitin-activating enzyme family. The encoded enzyme is a retinoid target that triggers promyelocytic leukemia (PML)/retinoic acid receptor alpha (RARalpha) degradation and apoptosis in acute promyelocytic leukemia, where it is involved in the conjugation of the ubiquitin-like interferon-stimulated gene 15 protein.</p>
<b>Applications</b>	Western Blotting: 1/500 - 1/2000. ELISA: Propose dilution 1/10000. Not yet tested in other applications. Determining optimal working dilutions by titration test.
<b>Limitations</b>	<ol style="list-style-type: none"><li>1. Conjugates with brighter fluorochromes, like PE and APC, will have a greater separation than those with dyes like FITC and CyQ. When populations overlap, the percentage of positive cells using a selected marker can be affected by the choice of fluorescent label.</li><li>2. Use of monoclonal antibodies in patient treatment can interfere with antigen target recognition by this reagent. This should be taken into account when samples are analyzed from patients treated in this fashion. IQ Products has not characterized the effect of the presence of therapeutic antibodies on the performance of this reagent.</li><li>3. Reagents can be used in different combinations, therefore laboratories need to become familiar performance characteristics of each antibody in relation with the combined markers in normal and abnormal samples.</li><li>4. Reagent data performance is based on EDTA-treated blood. Reagent performance can be affected by the use of other anticoagulants.</li></ol>



### Handling and Storage

Antibodies are supplied in 0.01 M sodium phosphate, 0.15 M NaCl; pH 7.3, 0.03% sodiumazide (NaN<sub>3</sub>) or as ascetic fluid containing 0.03% sodiumazide. Store the vials at 2-8 °C for a maximum of 2 weeks and store at -20°C for longer term storage. Monoclonal antibodies should be protected from prolonged exposure to light when conjugated with fluorochromes. Reagents are stable for the period shown on the vial label when stored properly.

### Warranty

Products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied, which extend beyond the description on the label of the product. IQ Products is not liable for property damage, personal injury, or economic loss caused by the product.

### Characterization

To ensure consistently high-quality reagents, each batch of monoclonal antibody is tested for conformance with characteristics of a standard reagent. Representative flow cytometric data is included in this data sheet.

### Warning

All products contain sodiumazide. This chemical is poisonous and hazardous. Handling should be done by trained staff only.

- 
- References**
1. Cancer Res. 1992 Mar 15;52(6):1536-41
  2. Proc Natl Acad Sci U S A. 1993 Jul 1;90(13):6071-5.
  3. Proc Natl Acad Sci U S A. 2002 Mar 19;99(6):3806-11.
- 

### Explanation of used symbols



REF



RUO

LOT



EC REP



Consult instructions for use

Catalogue number

Sufficient for

Caution, consult accompanying document

Keep away from (sun)light

Biological risks

Temperature limitation (°C)

For Research Use Only

Batch code

Use by yyyy-mm-dd

Manufacturer

Authorized Representative in the European Community

Conformité Européenne (European Conformity)

---

		<b>Label - tandem</b>	<b>Ex -max (nm)</b>	<b>Em -max (nm)</b>
P	PURE	purified material	-	-
F	FITC	FITC	488	519
R	R-PE	PE	488, 532	578
C	CyQ	PE-Cy5.18	488, 532	667
A	APC		595, 633, 635, 647	660
PC	PerCP		488, 532	678
PCC	PerCP-Cy5.5		488, 532	695

---



**IQ Products BV**

Rozenburglaan 13a  
9727 DL Groningen, The Netherlands

+31 (0)50 57 57 000

+31 (0)50 57 57 002

Technical [marketing@iqproducts.nl](mailto:marketing@iqproducts.nl)

Orders [orders@iqproducts.nl](mailto:orders@iqproducts.nl)

[www.iqproducts.nl](http://www.iqproducts.nl)