

## PRODUCT INFORMATION SHEET

### Mouse Monoclonal Antibody to ESR1

Format: PURE      RUO      REF      IQP-1464P      Volume: 0.1ml



#### Description

<b>Clone</b>	6B6
<b>Isotype</b>	Mouse IgG1
<b>Specificity</b>	ESR1
<b>Alternative names</b>	ER; ESR; Era; ESRA; NR3A1; DKFZp686N23123; ESR1
<b>Species</b>	Human
<b>Immunogen</b>	Purified recombinant fragment of human ESR1 expressed in E. Coli.
<b>Mw</b>	66kDa
<b>Format</b>	Ascitic fluid containing 0.03% sodium azide.

**Summary** This gene encodes an estrogen receptor, a ligand-activated transcription factor composed of several domains important for hormone binding, DNA binding, and activation of transcription. The protein localizes to the nucleus where it may form a homodimer or a heterodimer with estrogen receptor 2. Estrogen and its receptors are essential for sexual development and reproductive function, but also play a role in other tissues such as bone. Estrogen receptors are also involved in pathological processes including breast cancer, endometrial cancer, and osteoporosis.

**Applications** Western Blotting: 1/500 - 1/2000. Immunohistochemistry: 1/200 - 1/1000. Immunofluorescence: 1/200 - 1/1000. ELISA: Propose dilution 1/10000. Not yet tested in other applications. Determining optimal working dilutions by titration test.

#### Limitations

1. Conjugates with brighter fluorochromes, like PE and APC, will have a greater separation than those with dyes like FITC and CyQ. When populations overlap, the percentage of positive cells using a selected marker can be affected by the choice of fluorescent label.
2. Use of monoclonal antibodies in patient treatment can interfere with antigen target recognition by this reagent. This should be taken into account when samples are analyzed from patients treated in this fashion. IQ Products has not characterized the effect of the presence of therapeutic antibodies on the performance of this reagent.
3. Reagents can be used in different combinations, therefore laboratories need to become familiar performance characteristics of each antibody in relation with the combined markers in normal and abnormal samples.
4. Reagent data performance is based on EDTA-treated blood. Reagent performance can be affected by the use of other anticoagulants.



### Handling and Storage

Antibodies are supplied in 0.01 M sodium phosphate, 0.15 M NaCl; pH 7.3, 0.03% sodiumazide (NaN<sub>3</sub>) or as ascetic fluid containing 0.03% sodiumazide. Store the vials at 2-8 °C for a maximum of 2 weeks and store at -20°C for longer term storage. Monoclonal antibodies should be protected from prolonged exposure to light when conjugated with fluorochromes. Reagents are stable for the period shown on the vial label when stored properly.

### Warranty

Products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied, which extend beyond the description on the label of the product. IQ Products is not liable for property damage, personal injury, or economic loss caused by the product.

### Characterization

To ensure consistently high-quality reagents, each batch of monoclonal antibody is tested for conformance with characteristics of a standard reagent. Representative flow cytometric data is included in this data sheet.

### Warning

All products contain sodiumazide. This chemical is poisonous and hazardous. Handling should be done by trained staff only.

- References**
1. Calcif Tissue Int. 2010 Jul;87(1):25-35.
  2. Arch Gynecol Obstet. 2010 Jun 18.
  3. Biochemistry. 2010 Jul 27;49(29):5978-88.

### Explanation of used symbols



REF

Consult instructions for use

Catalogue number



Sufficient for



Caution, consult accompanying document



Keep away from (sun)light



Biological risks



Temperature limitation (°C)

RUO

For Research Use Only

LOT

Batch code



Use by yyyy-mm-dd



Manufacturer

EC/REP

Authorized Representative in the European Community



Conformité Européenne (European Conformity)

		Label - tandem	Ex -max (nm)	Em -max (nm)
P	PURE	purified material	-	-
F	FITC	FITC	488	519
R	R-PE	PE	488, 532	578
C	CyQ	PE-Cy5.18	488, 532	667
A	APC		595, 633, 635, 647	660
PC	PerCP		488, 532	678
PCC	PerCP-Cy5.5		488, 532	695



**IQ Products BV**

Rozenburglaan 13a  
9727 DL Groningen, The Netherlands

+31 (0)50 57 57 000

+31 (0)50 57 57 002

Technical [marketing@iqproducts.nl](mailto:marketing@iqproducts.nl)

Orders [orders@iqproducts.nl](mailto:orders@iqproducts.nl)

[www.iqproducts.nl](http://www.iqproducts.nl)