

## PRODUCT INFORMATION SHEET

### Mouse Monoclonal Antibody to IGF2BP3

Format: PURE      RUO      REF      IQP-1497P      Volume: 0.1ml

 Description	
<b>Clone</b>	8F11
<b>Isotype</b>	Mouse IgG1
<b>Specificity</b>	IGF2BP3
<b>Alternative names</b>	CT98; IMP3; KOC1; IMP-3; VICKZ3; DKFZp686F1078; IGF2BP3
<b>Species</b>	Human
<b>Immunogen</b>	Purified recombinant fragment of human IGF2BP3 expressed in E. Coli.
<b>Mw</b>	69kDa
<b>Format</b>	Ascitic fluid containing 0.03% sodium azide.
<b>Summary</b>	<p>The protein encoded by this gene is primarily found in the nucleolus, where it can bind to the 5' UTR of the insulin-like growth factor II leader 3 mRNA and may repress translation of insulin-like growth factor II during late development. The encoded protein contains several KH domains, which are important in RNA binding and are known to be involved in RNA synthesis and metabolism. Tissue specificity: Expressed in fetal liver, fetal lung, fetal kidney, fetal thymus, fetal placenta, fetal follicles of ovary and gonocytes of testis, growing oocytes, spermatogonia and semen (at protein level). Expressed in cervix adenocarcinoma, in testicular, pancreatic and renal-cell carcinomas (at protein level). Expressed ubiquitously during fetal development at 8 and 14 weeks of gestation. Expressed in ovary, testis, brain, placenta, pancreatic cancer tissues and pancreatic cancer cell lines. IMP-3 is a marker for carcinomas and high-grade dysplastic lesions of pancreatic ductal epithelium.</p>
<b>Applications</b>	Western Blotting: 1/500 - 1/2000. Immunohistochemistry: 1/200 - 1/1000. ELISA: Propose dilution 1/10000. Not yet tested in other applications. Determining optimal working dilutions by titration test.
<b>Limitations</b>	<ol style="list-style-type: none"><li>1. Conjugates with brighter fluorochromes, like PE and APC, will have a greater separation than those with dyes like FITC and CyQ. When populations overlap, the percentage of positive cells using a selected marker can be affected by the choice of fluorescent label.</li><li>2. Use of monoclonal antibodies in patient treatment can interfere with antigen target recognition by this reagent. This should be taken into account when samples are analyzed from patients treated in this fashion. IQ Products has not characterized the effect of the presence of therapeutic antibodies on the performance of this reagent.</li><li>3. Reagents can be used in different combinations, therefore laboratories need to become familiar performance characteristics of each antibody in relation with the combined markers in normal and abnormal samples.</li><li>4. Reagent data performance is based on EDTA-treated blood. Reagent performance can be affected by the use of other anticoagulants.</li></ol>



### Handling and Storage

Antibodies are supplied in 0.01 M sodium phosphate, 0.15 M NaCl; pH 7.3, 0.03% sodiumazide (NaN<sub>3</sub>) or as ascetic fluid containing 0.03% sodiumazide. Store the vials at 2-8 °C for a maximum of 2 weeks and store at -20°C for longer term storage. Monoclonal antibodies should be protected from prolonged exposure to light when conjugated with fluorochromes. Reagents are stable for the period shown on the vial label when stored properly.

### Warranty

Products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied, which extend beyond the description on the label of the product. IQ Products is not liable for property damage, personal injury, or economic loss caused by the product.

### Characterization

To ensure consistently high-quality reagents, each batch of monoclonal antibody is tested for conformance with characteristics of a standard reagent. Representative flow cytometric data is included in this data sheet.

### Warning

All products contain sodiumazide. This chemical is poisonous and hazardous. Handling should be done by trained staff only.

- 
- References** 1. Clin Cancer Res. 2008 Mar 15;14(6):1701-6.  
2. Hepatology. 2008 Oct;48(4):1118-27.
- 

### Explanation of used symbols



Consult instructions for use  
Catalogue number



Sufficient for



Caution, consult accompanying document



Keep away from (sun)light



Biological risks



Temperature limitation (°C)



For Research Use Only



Batch code



Use by yyyy-mm-dd



Manufacturer



Authorized Representative in the European Community



Conformité Européenne (European Conformity)




---

		<b>Label - tandem</b>	<b>Ex -max (nm)</b>	<b>Em -max (nm)</b>
P	PURE	purified material	-	-
F	FITC	FITC	488	519
R	R-PE	PE	488, 532	578
C	CyQ	PE-Cy5.18	488, 532	667
A	APC		595, 633, 635, 647	660
PC	PerCP		488, 532	678
PCC	PerCP-Cy5.5		488, 532	695

---



IQ Products BV  
Rozenburglaan 13a  
9727 DL Groningen, The Netherlands

- +31 (0)50 57 57 000
- +31 (0)50 57 57 002
- Technical [marketing@iqproducts.nl](mailto:marketing@iqproducts.nl)
- Orders [orders@iqproducts.nl](mailto:orders@iqproducts.nl)
- [www.iqproducts.nl](http://www.iqproducts.nl)