

PRODUCT INFORMATION SHEET

Mouse Monoclonal Antibody to SNAI1

Format: PURE RUO REF IQP-1544P Volume: 100µl

Description

Clone 6D2

Isotype Mouse IgG1
Specificity SNAI1

Alternative names

SNA; SNAH; SLUGH2; dJ710H13.1; SNAI1

Species Human

Immunogen Purified recombinant fragment of human SNAI1 expressed in E. Coli.

Mw 29kDa

Format Ascitic fluid containing 0.03% sodium azide.

Summary

Snail is a zinc-finger transcription factor that can repress E-cadherin transcription. Downregulation of E-cadherin is associated with epithelial-mesenchymal transition during embryonic development, a process also exploited by invasive cancer cells. Indeed, loss of E-cadherin expression is correlated with the invasive properties of some tumors and there is a considerable inverse correlation between Snail and E-cadherin mRNA levels in epithelial tumor cell lines. In addition, Snail blocks the cell cycle and confers resistance to cell death. Phosphorylation of Snail by GSK-3 and PAK1 regulates its stability, cellular localization and function. Tissue specificity: Expressed in a variety of tissues with the highest expression in kidney.

Applications ELISA: 1/10000; WB: 1/500 - 1/2000

Limitations

- 1. Conjugates with brighter fluorochromes, like PE and APC, will have a greater separation than those with dyes like FITC and CyQ. When populations overlap, the percentage of positive cells using a selected marker can be affected by the choice of fluorescent label.
- 2. Use of monoclonal antibodies in patient treatment can interfere with antigen target recognition by this reagent. This should be taken into account when samples are analyzed from patients treated in this fashion. IQ Products has not characterized the effect of the presence of therapeutic antibodies on the performance of this reagent.
- 3. Reagents can be used in different combinations, therefore laboratories need to become familiar performance characteristics of each antibody in relation with the combined markers in normal and abnormal samples.
- 4. Reagent data performance is based on EDTA-treated blood. Reagent performance can be affected by the use of other anticoagulants.

△ ♦ / * □

Handling and Storage

Store the vials at 2-8 °C for a maximum of 2 weeks and store at -20°C for longer term storage. Monoclonal antibodies should be protected from prolonged exposure to light when conjugated with fluorochromes. Reagents are stable for the period shown on the vial label when stored properly.

IQP-1544 – SNAI1 Version 2

Warranty

Products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied, which extend beyond the description on the label of the product. IQ Products is not liable for property damage, personal injury, or economic loss caused by the product.

Characterization

To ensure consistently high-quality reagents, each batch of monoclonal antibody is tested for conformance with characteristics of a standard reagent.

Warning

All products contain sodiumazide. This chemical is poisonous and hazardous. Handling should be done by trained staff only.

References

1.Exp Cell Res. 2008 Aug 1;314(13):2448-53. 2.Mol Cell Biol. 2008 Aug;28(15):4772-81.

Explanation of used symbols

	Consult instructions for use	
REF	Catalogue number	
E	Sufficient for	
\triangle	Caution, consult accompanying docum	ent
*	Keep away from (sun)light	
፟	Biological risks	
*	Temperature limitation (°C)	
RUO	For Research Use Only	The state of the s
LOT	Batch code	
Σ	Use by yyyy-mm-dd	
"	Manufacturer	

		Label - tandem	Ex -max (nm)	Em -max (nm)
Р	PURE	purified material	-	-
F	FITC	FITC	488	519
R	R-PE	PE PE	488, 532	578
С	CyQ	PE-Cy5.18	488, 532	667 CELICE
Α	APC		595, 633, 635, 647	660
PC	PerCP		488, 532	678
PCC	PerCP-Cy5.5		488, 532	695



IQ Products BV

Rozenburglaan 13a 9727 DL Groningen, The Netherlands

2 +31 (0)50 57 57 0003 +31 (0)50 57 57 002

Technical marketing@iqproducts.nlOrders orders@iqproducts.nl

<u>www.igproducts.nl</u>

IQP-1544 – SNAI1 Version 2