

PRODUCT INFORMATION SHEET

	Format: PURE	RUO	REF	IQP-1602P	Volume:	100µl		
Ĩ	Description							
Clone	5D10D11							
sotype	Mouse IgG2a							
Specificity	NFKB1							
Alternative names	p50; KBF1; p105; EBP-1; MGC54151; NFKB-p50; NFkappaB; NF-kappaB; NFKB-p105; NF-kappa-E							
Species	Human							
mmunogen	Purified recombinant fragment of human NFKB1 expressed in E. Coli.							
1w	50kDa/105kDa							
ormat	Ascitic fluid containing 0.03% sodium azide.							
Summary	This gene encodes a 105 kD protein which can undergo cotranslational processing by the 265 proteasome to produce a 50 kD protein. The 105 kD protein is a Rel protein-specific transcription inhibitor and the 50 kD protein is a DNA binding subunit of the NF-kappa-B (NFKB) protein complex. NFKB is a transcription regulator that is activated by various intra- and extra-cellula stimuli such as cytokines, oxidant-free radicals, ultraviolet irradiation, and bacterial or vira products. Activated NFKB translocates into the nucleus and stimulates the expression of generatively of biological functions. Inappropriate activation of NFKB has been associated with a number of inflammatory diseases while persistent inhibition of NFKB leads to inappropriate immune cell development or delayed cell growth. Two transcript variants encoding different isoforms have been found for this gene.							
Applications	ELISA: 1/10000; WE 1/400	3: 1/500 - 1/2000	; IHC: 1/2	00 - 1/1000; ICC	: 1/200 - 1/10	000; FCM: 1/200		
Limitations	 Conjugates with brighter fluorochromes, like PE and APC, will have a greater separation that those with dyes like FITC and CyQ. When populations overlap, the percentage of positive ceusing a selected marker can be affected by the choice of fluorescent label. Use of monoclonal antibodies in patient treatment can interfere with antigen target recognition by this reagent. This should be taken into account when samples are analyzed from patient treated in this fashion. IQ Products has not characterized the effect of the presence 							
	therapeutic antibodies on the performance of this reagent.3. Reagents can be used in different combinations, therefore laboratories need to be familiar performance characteristics of each antibody in relation with the combined marke							
	normal and abno	rmai sampies.				formance can b		

▲ ֎ ¥ 茶 🛛 Handling and Storage

Store the vials at 2-8 °C for a maximum of 2 weeks and store at -20°C for longer term storage. Monoclonal antibodies should be protected from prolonged exposure to light when conjugated with fluorochromes. Reagents are stable for the period shown on the vial label when stored properly.

Warranty	Products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied, which extend beyond the description on the label of the product. IQ Products is not liable for property damage, personal injury, or economic loss caused by the product.
Characteriz	ation To ensure consistently high-quality reagents, each batch of monoclonal antibody is tested for conformance with characteristics of a standard reagent.

Warning All products contain sodiumazide. This chemical is poisonous and hazardous. Handling should be done by trained staff only.

References

1.Cytokine. 2010 Feb;49(2):215-20. 2.Chemotherapy. 2009;55(5):381-5.

Explanation of used symbols

	Consult instructions	for use							
REF	Catalogue number								
$\overline{\mathbb{V}}$	Sufficient for								
$\mathbf{\Lambda}$	Caution, consult accompanying document								
*		Keep away from (sun)light							
- &	Biological risks								
×	-	ion (PC)							
RUO	Temperature limitat								
	For Research Use O	niy							
LOT	Batch code								
_ ≚	Use by yyyy-mm-do	ł							
- 	Manufacturer								
			rod	ucts					
		Label - tandem	Ex -max (nm)	Em -max (nm)					
Р	PURE	purified material		EIII - IIIax (IIIII)					
F	FITC	FITC	488	519					
R	R-PE	PE	488, 532						
C	CyQ	PE-Cy5.18	488, 532	Cence					
А	APC		595, 633, 635, 647	660					
PC	PerCP		488, 532	678					
PCC	PerCP-Cy5.5		488, 532	695					
	IQ Products BV								
	Rozenburglaan 13a								

9727 DL Groningen, The Netherlands