

#### PRODUCT INFORMATION SHEET

# **Mouse Monoclonal Antibody to BMI1**

Format: PURE RUO REF IQP-1653P Volume: 100µl

**Description** Ti

Clone 3E3

Mouse IgG1 **Isotype** 

**Specificity** 

**Alternative** names

PCGF4; RNF51; MGC12685; FLVI2/BMI1

**Species** Human

Immunogen Purified recombinant fragment of human BMI1 expressed in E. Coli.

Mw 37kDa

**Format** Ascitic fluid containing 0.03% sodium azide.

#### **Summary**

Component of the Polycomb group (PcG) multiprotein PRC1 complex, a complex required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1 complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility. In the PRC1 complex, it is required to stimulate the E3 ubiquitin-protein ligase activity of RNF2/RING2.

**Applications** 

ELISA: 1/10000; WB: 1/500 - 1/2000; IHC: 1/200 - 1/1000; ICC: 1/200 - 1/1000; FCM: 1/200 -

1/400

#### Limitations

- 1. Conjugates with brighter fluorochromes, like PE and APC, will have a greater separation than those with dyes like FITC and CyQ. When populations overlap, the percentage of positive cells using a selected marker can be affected by the choice of fluorescent label.
- 2. Use of monoclonal antibodies in patient treatment can interfere with antigen target recognition by this reagent. This should be taken into account when samples are analyzed from patients treated in this fashion. IQ Products has not characterized the effect of the presence of therapeutic antibodies on the performance of this reagent.
- 3. Reagents can be used in different combinations, therefore laboratories need to become familiar performance characteristics of each antibody in relation with the combined markers in normal and abnormal samples.
- 4. Reagent data performance is based on EDTA-treated blood. Reagent performance can be affected by the use of other anticoagulants.

# **△ ♦ / \*** □

#### **Handling and Storage**

Store the vials at 2-8 °C for a maximum of 2 weeks and store at -20°C for longer term storage. Monoclonal antibodies should be protected from prolonged exposure to light when conjugated with fluorochromes. Reagents are stable for the period shown on the vial label when stored properly.

# Warranty

Products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied, which extend beyond the description on the label of the product. IQ Products is not liable for property damage, personal injury, or economic loss caused by the product.

IQP-1653 - BMI1 Version 1

## Characterization

To ensure consistently high-quality reagents, each batch of monoclonal antibody is tested for conformance with characteristics of a standard reagent.

Warning

All products contain sodiumazide. This chemical is poisonous and hazardous. Handling should be done by trained staff only.

## References

1.Mol Cancer. 2009 Nov 10;8:98.

2.Cancer Res. 2009 Dec 1;69(23):9090-5.

## **Explanation of used symbols**

(li	Consult instructions for use
REF	Catalogue number
\$	Sufficient for
$\triangle$	Caution, consult accompanying document
*	Keep away from (sun)light
፟	Biological risks
*	Temperature limitation (°C)
RUO	For Research Use Only
LOT	Batch code
Σ	Use by yyyy-mm-dd
	Manufacturer

		Label - tandem	Ex -max (nm)	Em -max (nm)
P	PURE	purified material	14001	
F	FITC	FITC	488	519
R	R-PE	PE	488, 532	578
С	CyQ	PE-Cy5.18	488, 532	667
Α	APC		595, 633, 635, 647	660
PC	PerCP		488, 532	678
PCC	PerCP-Cy5.5	bric	488, 532	695 canca



IQ Products BV

Rozenburglaan 13a 9727 DL Groningen, The Netherlands

→ +31 (0)50 57 57 000

**431** (0)50 57 57 002

Technical Orders Technical marketing@igproducts.nl
Orders orders@igproducts.nl

<u>www.iqproducts.nl</u>

IQP-1653 - BMI1 Version 1