

PRODUCT INFORMATION SHEET

Mouse Monoclonal Antibody to TWIST1

Format: PURE RUO REF IQP-1794P Quantity: 0.1mg

Description

Clone 2F8E7

Isotype Mouse IgG2b
Specificity TWIST1

Alternative names

SCS; ACS3; CRS1; BPES2; BPES3; TWIST; bHLHa38

Species Human

Immunogen Purified recombinant fragment of human TWIST1 (AA: 9-74) expressed in E. Coli.

Mw 21kDa

Format Purified antibody in PBS with 0.05% sodium azide

Summary

Basic helix-loop-helix (bHLH) transcription factors have been implicated in cell lineage determination and differentiation. The protein encoded by this gene is a bHLH transcription factor and shares similarity with another bHLH transcription factor, Dermo1. The strongest expression of this mRNA is in placental tissue; in adults, mesodermally derived tissues express this mRNA preferentially. Mutations in this gene have been found in patients with Saethre-Chotzen syndrome.

Applications ELISA: 1/10000; WB: 1/500 - 1/2000

Limitations

- 1. Conjugates with brighter fluorochromes, like PE and APC, will have a greater separation than those with dyes like FITC and CyQ. When populations overlap, the percentage of positive cells using a selected marker can be affected by the choice of fluorescent label.
- 2. Use of monoclonal antibodies in patient treatment can interfere with antigen target recognition by this reagent. This should be taken into account when samples are analyzed from patients treated in this fashion. IQ Products has not characterized the effect of the presence of therapeutic antibodies on the performance of this reagent.
- 3. Reagents can be used in different combinations, therefore laboratories need to become familiar performance characteristics of each antibody in relation with the combined markers in normal and abnormal samples.
- 4. Reagent data performance is based on EDTA-treated blood. Reagent performance can be affected by the use of other anticoagulants.

⚠ ♣ **∦** ★ 🛂

Handling and Storage

Store the vials at 2-8 °C for a maximum of 2 weeks and store at -20°C for longer term storage. Monoclonal antibodies should be protected from prolonged exposure to light when conjugated with fluorochromes. Reagents are stable for the period shown on the vial label when stored properly.

Warranty

Products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied, which extend beyond the description on the label of the product. IQ Products is not liable for property damage, personal injury, or economic loss caused by the product.

IQP-1794 – TWIST1 Version 1

Characterization

To ensure consistently high-quality reagents, each batch of monoclonal antibody is tested for conformance with characteristics of a standard reagent.

Warning

All products contain sodiumazide. This chemical is poisonous and hazardous. Handling should be done by trained staff only.

References

1.J Cancer Res Clin Oncol. 2011 Oct;137(10):1487-93.

2.Cancer Res. 2011 Jan 1;71(1):245-54.

Explanation of used symbols



		Label - tandem	Ex -max (nm)	Em -max (nm)
Р	PURE	purified material		<u>-</u>
F	FITC	FITC	488	519
R	R-PE	PE	488, 532	578
C	CyQ	PE-Cy5.18	488, 532	667
Α	APC		595, 633, 635, 647	660
PC	PerCP		488, 532	678
PCC	PerCP-Cy5.5		488, 532	695
		brio	iht tluor	escence



IQ Products BV

Rozenburglaan 13a 9727 DL Groningen, The Netherlands

★ +31 (0)50 57 57 000★ +31 (0)50 57 57 002

Technical marketing@iqproducts.nl
 Orders orders@iqproducts.nl

www.iqproducts.nl

IQP-1794 – TWIST1 Version 1