

PRODUCT INFORMATION SHEET

Mouse Monoclonal Antibody to CD87

Format: PURE RUO REF IQP-1818P QTY 0.1mg

Description Ti

Clone 1D1B11

Mouse IgG2b **Isotype**

Specificity CD87

Alternative

PLAUR; UPAR; URKR; U-PAR names

Species Human

Immunogen Purified recombinant fragment of human CD87 (AA: 23-305) expressed in E. Coli.

37kDa Mw

Format Purified antibody in PBS with 0.05% sodium azide

Summary

This gene encodes the receptor for urokinase plasminogen activator and, given its role in localizing and promoting plasmin formation, likely influences many normal and pathological processes related to cell-surface plasminogen activation and localized degradation of the extracellular matrix. It binds both the proprotein and mature forms of urokinase plasmingen activator and permits the activation of the receptor-bound pro-enzyme by plasmin. The protein lacks transmembrane or cytoplasmic domains and may be anchored to the plasma membrane by a glycosylphosphatidylinositol (GPI) moiety following cleavage of the nascent polypeptide near its carboxyterminus. However, a soluble protein is also produced in some cell types. Alternative splicing results in multiple transcript variants encoding different isoforms. The proprotein experiences several post-translational cleavage reactions that have not yet been fully defined.

Applications ELISA: 1/10000; WB: 1/500 - 1/2000; FCM: 1/200 - 1/400; ICC: N/A; IHC: N/A.

Limitations

- 1. Conjugates with brighter fluorochromes, like PE and APC, will have a greater separation than those with dyes like FITC and CyQ. When populations overlap, the percentage of positive cells using a selected marker can be affected by the choice of fluorescent label.
- 2. Use of monoclonal antibodies in patient treatment can interfere with antigen target recognition by this reagent. This should be taken into account when samples are analyzed from patients treated in this fashion. IQ Products has not characterized the effect of the presence of therapeutic antibodies on the performance of this reagent.
- 3. Reagents can be used in different combinations, therefore laboratories need to become familiar performance characteristics of each antibody in relation with the combined markers in normal and abnormal samples.
- 4. Reagent data performance is based on EDTA-treated blood. Reagent performance can be affected by the use of other anticoagulants.

△ ♦ ∤ * □

Handling and Storage

Store the vials at 2-8°C for a maximum of 2 weeks and store at -20°C for longer term storage. Monoclonal antibodies should be protected from prolonged exposure to light when conjugated with fluorochromes. Reagents are stable for the period shown on the vial label when stored properly.

Warranty

Products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied,

IQP-1818 - CD87 Version 2 which extend beyond the description on the label of the product. IQ Products is not liable for property damage, personal injury, or economic loss caused by the product.

Characterization

To ensure consistently high-quality reagents, each batch of monoclonal antibody is tested for conformance with characteristics of a standard reagent.

Warning

All products contain sodium azide. This chemical is poisonous and hazardous. Handling should be done by trained staff only.

References

1.Biomark Med. 2016;10(2):209-16. 2.Andrology. 2015 Nov;3(6):1054-61.

Explanation of used symbols



D	DUDE	Label - tandem	Ex -max (nm)	Em -max (nm)
P	PURE	purified material		
F	FITC	FITC	488	519
R	R-PE	PE	488, 532	578
С	CyQ	PE-Cy5.18	488, 532	667
Α	APC	brio	595, 633, 635, 647	660
PC	PerCP		488, 532	678
PCC	PerCP-Cy5.5		488, 532	695



IQ Products BV

Rozenburglaan 13a 9727 DL Groningen, The Netherlands

431 (0)50 57 57 002

 $\begin{tabular}{ll} \hline @ Technical & \underline{marketing@iqproducts.nl} \\ \hline \end{tabular}$

Orders orders@igproducts.nl

<u>www.igproducts.nl</u>

IQP-1818 – CD87 Version 2