

### **PRODUCT INFORMATION SHEET**

## **Mouse Monoclonal Antibody to MSLN**

Format: PURE RUO REF IQP-1823P QTY 0.1mg

Description

Clone 5C11A4

Isotype Mouse IgG1

**Specificity** MSLN

Alternative names

MPF; SMRP

Species Human

Immunogen Purified recombinant fragment of human MSLN (AA: 37-286) expressed in E. Coli.

Mw 69kDa

**Format** Purified antibody in PBS with 0.05% sodium azide

#### Summary

This gene encodes a preproprotein that is proteolytically processed to generate two protein products, megakaryocyte potentiating factor and mesothelin. Megakaryocyte potentiating factor functions as a cytokine that can stimulate colony formation of bone marrow megakaryocytes. Mesothelin is a glycosylphosphatidylinositol-anchored cell-surface protein that may function as a cell adhesion protein. This protein is overexpressed in epithelial mesotheliomas, ovarian cancers and in specific squamous cell carcinomas. Alternative splicing results in multiple transcript variants, at least one of which encodes an isoform that is proteolytically processed.

**Applications** WB:1/500 - 1/2000; FCM:1/200 - 1/400; ELISA:1/10000.

### Limitations

- 1. Conjugates with brighter fluorochromes, like PE and APC, will have a greater separation than those with dyes like FITC and CyQ. When populations overlap, the percentage of positive cells using a selected marker can be affected by the choice of fluorescent label.
- Use of monoclonal antibodies in patient treatment can interfere with antigen target recognition by this reagent. This should be taken into account when samples are analyzed from patients treated in this fashion. IQ Products has not characterized the effect of the presence of therapeutic antibodies on the performance of this reagent.
- 3. Reagents can be used in different combinations, therefore laboratories need to become familiar performance characteristics of each antibody in relation with the combined markers in normal and abnormal samples.
- 4. Reagent data performance is based on EDTA-treated blood. Reagent performance can be affected by the use of other anticoagulants.

## **▲ ♦ / \* ×**

## **Handling and Storage**

Store the vials at  $2-8^{\circ}$ C for a maximum of 2 weeks and store at  $-20^{\circ}$ C for longer term storage. Monoclonal antibodies should be protected from prolonged exposure to light when conjugated with fluorochromes. Reagents are stable for the period shown on the vial label when stored properly.

# Warranty

Products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied, which extend beyond the description on the label of the product. IQ Products is not liable for property damage, personal injury, or economic loss caused by the product.

IQP-1823 – MSLN Version 2

## Characterization

To ensure consistently high-quality reagents, each batch of monoclonal antibody is tested for conformance with characteristics of a standard reagent.

**Warning** All products contain sodium azide. This chemical is poisonous and hazardous. Handling should be done by trained staff only.

### References

1.Tumour Biol. 2018 Oct;40(10):1010428318804937.

2.Mol Cancer. 2017 Mar 14;16(1):63.

## **Explanation of used symbols**



		Label - tandem	Ex -max (nm)	Em -max (nm)
P	PURE	purified material		- ·
F	FITC	FITC	488	519
R	R-PE	PE	488, 532	578
C	CyQ	PE-Cy5.18	488, 532	667
Α	APC		595, 633, 635, 647	660
PC	PerCP		488, 532	678
PCC	PerCP-Cy5.5		488, 532	695
		brio	ibt tluor	OCCODCO



IQ Products BV

Rozenburglaan 13a 9727 DL Groningen, The Netherlands

**+31** (0)50 57 57 002

Technical marketing@iqproducts.nlOrders orders@iqproducts.nl

www.iqproducts.nl

IQP-1823 – MSLN Version 2