

#### **PRODUCT INFORMATION SHEET**

# **Mouse Monoclonal Antibody to CD147**

Format: PURE RUO REF IQP-1840P QTY 0.1mg

Description

Clone 3E10C1
Isotype Mouse IgG1
Specificity CD147

Alternative names

BSG; OK; 5F7; TCSF; EMMPRIN

Species Human

Immunogen Purified recombinant fragment of human CD147 (AA: extra 138-323) expressed in E. Coli.

**Mw** 42.2kDa

**Format** Purified antibody in PBS with 0.05% sodium azide

Summary The protein encoded by this gene is a plasma membrane protein that is important in

spermatogenesis, embryo implantation, neural network formation, and tumor progression. The encoded protein is also a member of the immunoglobulin superfamily. Multiple transcript variants

encoding different isoforms have been found for this gene.

**Applications** ELISA: 1/10000; WB: 1/500 - 1/2000; ICC: N/A; FCM: 1/200 - 1/400; IHC: N/A.

### Limitations

- 1. Conjugates with brighter fluorochromes, like PE and APC, will have a greater separation than those with dyes like FITC and CyQ. When populations overlap, the percentage of positive cells using a selected marker can be affected by the choice of fluorescent label.
- 2. Use of monoclonal antibodies in patient treatment can interfere with antigen target recognition by this reagent. This should be taken into account when samples are analyzed from patients treated in this fashion. IQ Products has not characterized the effect of the presence of therapeutic antibodies on the performance of this reagent.
- 3. Reagents can be used in different combinations, therefore laboratories need to become familiar performance characteristics of each antibody in relation with the combined markers in normal and abnormal samples.
- 4. Reagent data performance is based on EDTA-treated blood. Reagent performance can be affected by the use of other anticoagulants.

# **△ ♦ / \* ×**

#### **Handling and Storage**

Store the vials at 2-8°C for a maximum of 2 weeks and store at -20°C for longer term storage. Monoclonal antibodies should be protected from prolonged exposure to light when conjugated with fluorochromes. Reagents are stable for the period shown on the vial label when stored properly.

### Warranty

Products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied, which extend beyond the description on the label of the product. IQ Products is not liable for property damage, personal injury, or economic loss caused by the product.

# Characterization

To ensure consistently high-quality reagents, each batch of monoclonal antibody is tested for conformance with characteristics of a standard reagent.

IQP-1840 - CD147 Version 2

Warning All products contain sodium azide. This chemical is poisonous and hazardous. Handling should be done by trained staff only.

#### References

1.Oncotarget. 2016 Feb 2;7(5):5613-29. 2.J Biochem. 2016 May;159(5):481-90.

## **Explanation of used symbols**

(Ii Consult instructions for use REF Catalogue number Sufficient for  $\triangle$ Caution, consult accompanying document \* Keep away from (sun)light **₽** Biological risks Temperature limitation (°C) RUO For Research Use Only LOT Batch code Use by yyyy-mm-dd Manufacturer

		Label - tandem	Ex -max (nm)	Em -max (nm)
Р	PURE	purified material	-	-
F	FITC	FITC	488	519
R	R-PE	PE	488, 532	578
C	CyQ	PE-Cy5.18	488, 532	667
Α	APC		595, 633, 635, 647	660
PC	PerCP		488, 532	678
PCC	PerCP-Cy5.5		488, 532	695
	·		•	

right fluorescence

**IQ Products BV** 

Rozenburglaan 13a 9727 DL Groningen, The Netherlands

+31 (0)50 57 57 000

+31 (0)50 57 57 002

Technical marketing@iqproducts.nl

Orders orders@iqproducts.nl

www.iqproducts.nl

IQP-1840 - CD147 Version 2