

PRODUCT INFORMATION SHEET

Mouse Monoclonal Antibody to PCNA								
	Format: PURE	RUO	REF	IQP-1849P	Volume:	100µl		
	Description							
Clone	7H4F8							
Isotype	Mouse IgG1							
Specificity	PCNA							
Alternative names								
Species	Human, Monkey							
Immunogen	Purified recombinant fragment of human PCNA (AA: 53-196) expressed in E. Coli.							
Mw	28.7kDa							
Format	Ascitic fluid containing 0.03% sodium azide.							
Summary	The protein encoded by this gene is found in the nucleus and is a cofactor of DNA polymerase delta. The encoded protein acts as a homotrimer and helps increase the processivity of leading strand synthesis during DNA replication. In response to DNA damage, this protein is ubiquitinated and is involved in the RAD6-dependent DNA repair pathway. Two transcript variants encoding the same protein have been found for this gene. Pseudogenes of this gene have been described on chromosome 4 and on the X chromosome.							
Applications	ELISA: 1/10000; WB	: 1/500 - 1/2000;	IHC: 1/20	0 - 1/1000; FCM:	1/200 - 1/400			

Limitations

- 1. Conjugates with brighter fluorochromes, like PE and APC, will have a greater separation than those with dyes like FITC and CyQ. When populations overlap, the percentage of positive cells using a selected marker can be affected by the choice of fluorescent label.
 - 2. Use of monoclonal antibodies in patient treatment can interfere with antigen target recognition by this reagent. This should be taken into account when samples are analyzed from patients treated in this fashion. IQ Products has not characterized the effect of the presence of therapeutic antibodies on the performance of this reagent.
 - 3. Reagents can be used in different combinations, therefore laboratories need to become familiar performance characteristics of each antibody in relation with the combined markers in normal and abnormal samples.
 - 4. Reagent data performance is based on EDTA-treated blood. Reagent performance can be affected by the use of other anticoagulants.

▲ ֎ ¥ 苯 ¥ Handling and Storage

Store the vials at 2-8 °C for a maximum of 2 weeks and store at -20°C for longer term storage. Monoclonal antibodies should be protected from prolonged exposure to light when conjugated with fluorochromes. Reagents are stable for the period shown on the vial label when stored properly.

Warranty	Products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied, which extend beyond the description on the label of the product. IQ Products is not liable for property damage, personal injury, or economic loss caused by the product.
Characteriza	tion

Characterization

To ensure consistently high-quality reagents, each batch of monoclonal antibody is tested for conformance with characteristics of a standard reagent.

Warning All products contain sodiumazide. This chemical is poisonous and hazardous. Handling should be done by trained staff only.

References

1.Cancer Res. 2012 Jul 1;72(13):3217-27. 2.PLoS One. 2012;7(1):e29416.

Explanation of used symbols

REF ♥ ▲ ** @ LOT ₽	Consult instructions Catalogue number Sufficient for Caution, consult acc Keep away from (su Biological risks Temperature limitat For Research Use On Batch code Use by yyyy-mm-dd Manufacturer	ompanying document n)light ion (°C) nly		ucts
P F R C A PC PCC	PURE FITC R-PE CyQ APC PerCP PerCP-Cy5.5	Label - tandem purified material FITC PE PE-Cy5.18	Ex -max (nm) - 488 488, 532 488, 532 595, 633, 635, 647 488, 532 488, 532	Em -max (nm) - 519 578 667 660 678 695
	Rozenburglaan 13a 9727 DL Groningen, T	he Netherlands		

+31 (0)50 57 57 000

- ∃ +31 (0)50 57 57 002
- Image: Marketing@iqproducts.nl
 marketing@iqproducts.nl

 Image: Marketing@iqproducts.nl
 orders@iqproducts.nl
- <u>www.iqproducts.nl</u>