

PRODUCT INFORMATION SHEET

Mouse Monoclonal Antibody to CD152

Format: PURE RUO REF IQP-1852P Quantity: 0.1mg

Description

Clone 8B3F8

Isotype Mouse IgG1
Specificity CD152

Alternative names

CTLA4; CD; GSE; GRD4; ALPS5; CTLA-4; IDDM12; CELIAC3

Species Human

Immunogen Purified recombinant fragment of human CD152 (AA: extra 36-161) expressed in E. Coli.

Mw 24.6kDa

Format Purified antibody in PBS with 0.05% sodium azide

Summary

This gene is a member of the immunoglobulin superfamily and encodes a protein which transmits an inhibitory signal to T cells. The protein contains a V domain, a transmembrane domain, and a cytoplasmic tail. Alternate transcriptional splice variants, encoding different isoforms, have been characterized. The membrane-bound isoform functions as a homodimer interconnected by a disulfide bond, while the soluble isoform functions as a monomer. Mutations in this gene have been associated with insulin-dependent diabetes mellitus, Graves disease, Hashimoto thyroiditis, celiac disease, systemic lupus erythematosus, thyroid-associated orbitopathy, and other autoimmune diseases.

Applications ELISA: 1/10000; WB: 1/500 - 1/2000; ICC: N/A; FCM: 1/200 - 1/400; IHC: N/A

Limitations

- 1. Conjugates with brighter fluorochromes, like PE and APC, will have a greater separation than those with dyes like FITC and CyQ. When populations overlap, the percentage of positive cells using a selected marker can be affected by the choice of fluorescent label.
- 2. Use of monoclonal antibodies in patient treatment can interfere with antigen target recognition by this reagent. This should be taken into account when samples are analyzed from patients treated in this fashion. IQ Products has not characterized the effect of the presence of therapeutic antibodies on the performance of this reagent.
- 3. Reagents can be used in different combinations, therefore laboratories need to become familiar performance characteristics of each antibody in relation with the combined markers in normal and abnormal samples.
- 4. Reagent data performance is based on EDTA-treated blood. Reagent performance can be affected by the use of other anticoagulants.

Handling and Storage

Store the vials at 2-8 °C for a maximum of 2 weeks and store at -20°C for longer term storage. Monoclonal antibodies should be protected from prolonged exposure to light when conjugated with fluorochromes. Reagents are stable for the period shown on the vial label when stored properly.

Warranty

Products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied, which extend beyond the description on the label of the product. IQ Products is not liable for property damage, personal injury, or economic loss caused by the product.

IQP-1852 - CD152 Version 1

Characterization

To ensure consistently high-quality reagents, each batch of monoclonal antibody is tested for conformance with characteristics of a standard reagent.

Warning

All products contain sodiumazide. This chemical is poisonous and hazardous. Handling should be done by trained staff only.

References

1.Asian Pac J Cancer Prev. 2016;17(8):3785-91. 2.Eur J Cancer. 2015 Nov;51(17):2689-97.

Explanation of used symbols



		Label - tandem	Ex -max (nm)	Em -max (nm)
P	PURE	purified material		<u>-</u>
F	FITC	FITC	488	519
R	R-PE	PE	488, 532	578
C	CyQ	PE-Cy5.18	488, 532	667
Α	APC		595, 633, 635, 647	660
PC	PerCP		488, 532	678
PCC	PerCP-Cy5.5		488, 532	695
		brid	iht tluor	escence



IQ Products BV

Rozenburglaan 13a 9727 DL Groningen, The Netherlands

= +31 (0)50 57 57 002

Technical <u>marketing@iqproducts.nl</u>
Orders <u>orders@iqproducts.nl</u>

www.iqproducts.nl

IQP-1852 - CD152 Version 1