

PRODUCT INFORMATION SHEET

Mouse Monoclonal Antibody to EGFR						
	Format: PURE	RUO	REF	IQP-1853P	Quantity:	0.1mg
	Description					
Clone	5E10D3					
Isotype	Mouse IgG2b					
Specificity	EGFR					
Alternative names	ERBB; HER1; mENA; ERBB1; PIG61					
Species	Human					
Immunogen	Purified recombinant fragment of human EGFR (AA: 693-893) expressed in E. Coli.					
Mw	175kDa					
Format	Purified antibody in PBS with 0.05% sodium azide					
Summary	The protein encoded by this gene is a transmembrane glycoprotein that is a member of the protein kinase superfamily. This protein is a receptor for members of the epidermal growth factor family EGFR is a cell surface protein that binds to epidermal growth factor. Binding of the protein to a ligand induces receptor dimerization and tyrosine autophosphorylation and leads to cell proliferation. Mutations in this gene are associated with lung cancer. Multiple alternatively spliced transcript variants that encode different protein isoforms have been found for this gene.					

Applications ELISA: 1/10000; WB: 1/500 - 1/2000; FCM: 1/200 - 1/400

Limitations

- 1. Conjugates with brighter fluorochromes, like PE and APC, will have a greater separation than those with dyes like FITC and CyQ. When populations overlap, the percentage of positive cells using a selected marker can be affected by the choice of fluorescent label.
- 2. Use of monoclonal antibodies in patient treatment can interfere with antigen target recognition by this reagent. This should be taken into account when samples are analyzed from patients treated in this fashion. IQ Products has not characterized the effect of the presence of therapeutic antibodies on the performance of this reagent.
- 3. Reagents can be used in different combinations, therefore laboratories need to become familiar performance characteristics of each antibody in relation with the combined markers in normal and abnormal samples.
- 4. Reagent data performance is based on EDTA-treated blood. Reagent performance can be affected by the use of other anticoagulants.

⚠ & ∦ 🕷 🎴

Handling and Storage

Store the vials at 2-8 °C for a maximum of 2 weeks and store at -20°C for longer term storage. Monoclonal antibodies should be protected from prolonged exposure to light when conjugated with fluorochromes. Reagents are stable for the period shown on the vial label when stored properly.

Warranty Products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied, which extend beyond the description on the label of the product. IQ Products is not liable for property damage, personal injury, or economic loss caused by the product.

Characterization

To ensure consistently high-quality reagents, each batch of monoclonal antibody is tested for conformance with characteristics of a standard reagent.

Warning

All products contain sodiumazide. This chemical is poisonous and hazardous. Handling should be done by trained staff only.

References

1.J Immunol. 2012 Dec 1;189(11):5230-9. 2.J Biol Chem. 2012 Oct 12;287(42):35201-11.

Explanation of used symbols



IQ Products BV

Rozenburglaan 13a

- 1 +31 (0)50 57 57 000
- +31 (0)50 57 57 002
- marketing@iqproducts.nl orders@iqproducts.nl M Technical
- Orders
- www.iqproducts.nl

⁹⁷²⁷ DL Groningen, The Netherlands