

## PRODUCT INFORMATION SHEET

# **Mouse Monoclonal Antibody to ABCG2**

Format: PURE RUO REF IQP-1856P Volume: 100µl

Description

Clone 3G8

Isotype Mouse IgG1
Specificity ABCG2

Alternative MRX; MXR; ABCP; BCRP; BMDP; MXR1; ABC15; BCRP1; CD338; CDw338; EST157481;

names MGC102821

**Species** Human, Mouse, Monkey

Immunogen Purified recombinant fragment of human ABCG2 expressed in E. Coli.

Mw 72kDa

**Format** Ascitic fluid containing 0.03% sodium azide.

## **Summary**

The membrane-associated protein encoded by this gene is included in the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the White subfamily. Alternatively referred to as a breast cancer resistance protein, this protein functions as a xenobiotic transporter which may play a major role in multi-drug resistance. It likely serves as a cellular defense mechanism in response to mitoxantrone and anthracycline exposure. Significant expression of this protein has been observed in the placenta, which may suggest a potential role for this molecule in placenta tissue. Tissue specificity: Highly expressed in placenta. Low expression in small intestine, liver and colon.

Applications ELISA: 1/10000; WB: 1/500 - 1/2000; IHC: 1/200 - 1/1000; ICC: 1/200 - 1/1000; FCM: 1/200 - 1/400

## Limitations

- 1. Conjugates with brighter fluorochromes, like PE and APC, will have a greater separation than those with dyes like FITC and CyQ. When populations overlap, the percentage of positive cells using a selected marker can be affected by the choice of fluorescent label.
- Use of monoclonal antibodies in patient treatment can interfere with antigen target recognition by this reagent. This should be taken into account when samples are analyzed from patients treated in this fashion. IQ Products has not characterized the effect of the presence of therapeutic antibodies on the performance of this reagent.
- 3. Reagents can be used in different combinations, therefore laboratories need to become familiar performance characteristics of each antibody in relation with the combined markers in normal and abnormal samples.
- 4. Reagent data performance is based on EDTA-treated blood. Reagent performance can be affected by the use of other anticoagulants.

# **△ ♦ / \* 2**

## **Handling and Storage**

Store the vials at 2-8 °C for a maximum of 2 weeks and store at -20°C for longer term storage. Monoclonal antibodies should be protected from prolonged exposure to light when conjugated with fluorochromes. Reagents are stable for the period shown on the vial label when stored properly.

IQP-1856 – ABCG2 Version 1

## Warranty

Products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied, which extend beyond the description on the label of the product. IQ Products is not liable for property damage, personal injury, or economic loss caused by the product.

## Characterization

To ensure consistently high-quality reagents, each batch of monoclonal antibody is tested for conformance with characteristics of a standard reagent.

# Warning

All products contain sodiumazide. This chemical is poisonous and hazardous. Handling should be done by trained staff only.

#### References

1.Carcinogenesis. 2008 Dec;29(12):2289-97.

2.Pharm Res. 2009 Feb;26(2):449-58.

# **Explanation of used symbols**



		Label - tandem	Ex -max (nm)	Em -max (nm)
P	PURE	purified material	-	-
F	FITC	FITC	488	519
R	R-PE	PE PE	488, 532	578
С	CyQ	PE-Cy5.18	488, 532	667
Α	APC	_	595, 633, 635, 647	660
PC	PerCP		488, 532	678
PCC	PerCP-Cy5.5		488, 532	695



IQ Products BV

Rozenburglaan 13a 9727 DL Groningen, The Netherlands

★ +31 (0)50 57 57 000★ +31 (0)50 57 57 002

Technical marketing@iqproducts.nl
Orders orders@iqproducts.nl

<u>www.igproducts.nl</u>

IQP-1856 – ABCG2 Version 1