

PRODUCT INFORMATION SHEET

Mouse Monoclonal Antibody to MUC5AC							
	Format: PURE RUO REF IQP-1872P Quantity: 0.1mg						
	Description						
Clone	7E9D8						
Isotype	Mouse IgG2b						
Specificity	MUC5AC						
Alternative names	TBM; leB; MUC5; mucin						
Species	Human						
Immunogen	Purified recombinant fragment of human MUC5AC (AA: 5528-5627) expressed in E. Coli.						
Mw	585kDa						
Format	Purified antibody in PBS with 0.05% sodium azide						
Summary	MUC5AC (Mucin 5AC, Oligomeric Mucus/Gel-Forming) is a Protein Coding gene. Diseases associated with MUC5AC include stomach cancer and pancreatic ductal carcinoma. Among its related pathways are O-linked glycosylation and Transport to the Golgi and subsequent modification. GO annotations related to this gene include extracellular matrix structural constituent. An important paralog of this gene is OTOG.						
Applications	ELISA: 1/10000; WB: 1/500 - 1/2000; FCM: 1/200 - 1/400						
Limitations	1. Conjugates with brighter fluorochromes, like PE and APC, will have a greater separation that						
	those with dyes like FITC and CyQ. When populations overlap, the percentage of positive ce						

- those with dyes like FITC and CyQ. When populations overlap, the percentage of positive cells using a selected marker can be affected by the choice of fluorescent label.Use of monoclonal antibodies in patient treatment can interfere with antigen target recognition
- by this reagent. This should be taken into account when samples are analyzed from patients treated in this fashion. IQ Products has not characterized the effect of the presence of therapeutic antibodies on the performance of this reagent.
- 3. Reagents can be used in different combinations, therefore laboratories need to become familiar performance characteristics of each antibody in relation with the combined markers in normal and abnormal samples.
- 4. Reagent data performance is based on EDTA-treated blood. Reagent performance can be affected by the use of other anticoagulants.

▲ ֎ ∦ 巻 🖁 Handling and Storage

Store the vials at 2-8 °C for a maximum of 2 weeks and store at -20°C for longer term storage. Monoclonal antibodies should be protected from prolonged exposure to light when conjugated with fluorochromes. Reagents are stable for the period shown on the vial label when stored properly.

Warranty Products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied, which extend beyond the description on the label of the product. IQ Products is not liable for property damage, personal injury, or economic loss caused by the product.

Characterization

To ensure consistently high-quality reagents, each batch of monoclonal antibody is tested for conformance with characteristics of a standard reagent.

All products contain sodiumazide. This chemical is poisonous and hazardous. Handling should be Warning done by trained staff only.

References

1.Biochemistry. 2015 Feb 3;54(4):1089-99. 2.Int J Oncol. 2013 Mar;42(3):887-93.

Explanation of used symbols

	Consult instructions for use				
REF	Catalogue number				
$\overline{\mathbb{V}}$	Sufficient for				
\wedge	Caution, consult accompanying document				
*	Keep away from (sun)light				
\$	Biological risks				
X	Temperature limitation (°C)				
RUO	For Research Use Only				
LOT	Batch code				
2	Use by yyyy-mm-dd				
	Manufacturer				

		Label - tandem	Ex -max (nm)	Em -max (nm)
Р	PURE	purified material		
F	FITC	FITC	488	519
R	R-PE	PE	488, 532	578
С	CyQ	PE-Cy5.18	488, 532	667
Α	APC		595, 633, 635, 647	660
PC	PerCP		488, 532	678
PCC	PerCP-Cy5.5	brio	488, 532	695 CODCO
				COLCILC

IQ Products BV

Rozenburglaan 13a 9727 DL Groningen, The Netherlands

1 +31 (0)50 57 57 000

₼ +31 (0)50 57 57 002

- ∕≞ ∕≏
 Technical
 marketing@iqproducts.nl

 Orders
 orders@iqproducts.nl

<u>www.iqproducts.nl</u>