

PRODUCT INFORMATION SHEET
 Monoclonal antibodies detecting human antigens

CD5 FITC **CD20** R-PE RUO REF IQP-210FR 50 tests

RUO *For Research Use Only*



Description

CD5 is present on all mature T cells and most thymocytes. It is also expressed on a small subpopulation of normal B cells. The majority of T-cell malignancies (76%) express CD5, and almost 85% of T-cell acute lymphoblastic leukemias are CD5 positive (1). Further, CD5 is expressed in some B-cell derived lymphoproliferative disorders, notably chronic lymphocytic leukemia (CLL) (> 90%), and centrocytic leukemia. CD20 can be considered a pan-B cell antigen, as it is expressed on the surface of all mature B lymphocytes but not in secreting plasma cells. CD20 is expressed early during pre-B cell development, presumably just before the expression of cytoplasmic μ -chains, and it persists until plasma cell differentiation. A small subset of T cells expresses low levels of CD20. CD20 is expressed in acute lymphoblastic leukemia, B-cell chronic lymphocytic leukemia, hairy cell leukemia and Burkitt's lymphoma. In order to perform appropriate analysis. Multicolor reagents are preferable to single-color reagents for the comprehensive analysis of flow cytometry specimens. The correct use of color compensation is particularly important in multicolor analysis.

CD5

Clone MCD5

Isotype Murine IgG2b

Specificity CD5 is present on all mature T cells and 95% of thymocytes. Anti-CD5 also reacts with a distinct subset of normal B cells and most CLL cells, but with few other B cell leukemias and lymphomas. CD5 appears to be a relatively late marker during B cell differentiation. CD5 expression is thought to be absent on surface Ig negative B-lineage cells but appears on IgM+ cells in both fetal liver and bone marrow

CD20

Clone B-ly1

Isotype Murine IgG1

Specificity Monoclonal antibodies clustered as CD19 detect all peripheral blood B cells. In addition, CD19 is expressed on precursor B cells during maturation, but not on mature plasma cells. The function of the CD19 molecule is related to signal transfer and is involved in regulation of B cell proliferation. CD19 is considered to be a characteristic B cell marker and therefore commonly used in routine immunophenotyping. CD19 may also be expressed on follicular dendritic cells.

HLDA Workshop


4th International Workshop on Human Leukocyte Differentiation Antigens.

Usage

All these reagents are effectively formulated for direct immunofluorescent staining of human tissue for flow cytometric analysis using 10 μ L/10⁶ leukocytes for singles and 20 μ L/10⁶ leukocytes in case of dual and triple combinations. Since applications vary, each investigator should titrate the reagent to obtain optimal results.

Representative Data

IQP-210FR (CD5/CD20) was analyzed by flow cytometry using a blood sample from a healthy volunteer. Direct staining was performed by adding 20 μ l of this dual to 100 μ l blood sample.



Immunofluorescence staining and lysing protocol

Flow cytometry method for use with dual and triple combinations

1. Add 100 µl of EDTA-treated blood (i.e. approx. 10^6 leukocytes) to a 5 ml reagent tube. The content of one tube is sufficient to perform one test.
For combinations with anti-kappa and/or anti-lambda Ig see application note below.
2. Add to each tube 20 µl of labeled monoclonal antibody combination*.
3. Vortex the tube to ensure thorough mixing of antibody and cells.
4. Incubate the tube for 15 minutes at room temperature in the dark.
5. Add 100 µl of IQ Lyse (IQP-199 ready-to-use) and mix immediately.
6. Incubate for 10 minutes at room temperature in the dark.
7. Add 2 ml of demineralized water and incubate for 10 minutes in the dark.
8. Centrifuge the labeled cell suspension for 2 minutes at 1000 x g.
9. Remove the supernatant and resuspend the cells in 200 µl of PBS**.
10. Analyze by flow cytometry within four hours (alternatively, the cells may be fixed by 0.05% of formaline in buffered saline for analysis the next day. Some antigens are readily destroyed upon fixation and this should be taken into account when using this alternative).

* Appropriate mouse Ig isotype control samples can be included in any labeling study

** PBS: Phosphate Buffered Saline, pH 7.2

Application note for anti-kappa and/or anti-lambda Ig combinations

Add 2 ml of PBS containing 0.001% (v/v) Heparin (**prewarmed to 37 °C**) to the cell suspension
Vortex, centrifuge (2 min at 300x g) and discard the supernatant
Repeat this step twice
Resuspend the pelleted blood cells in 100 µl PBS containing 0.001% (v/v) Heparin



Handling and Storage

Antibodies are supplied either as 100 tests per vial (1 ml) for singles or 50 tests per vial (1 ml) for dual and triple combinations. They are supplied in 0.01 M sodium phosphate, 0.15 M NaCl; pH 7.3, 0.2% BSA, 0.09% sodiumazide (NaN_3). Store the vials at 2-8 °C. Monoclonal antibodies should be protected from prolonged exposure to light. Reagents are stable for the period shown on the vial label when stored properly.

Warranty Products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied, which extend beyond the description on the label of the product. IQ Products is not liable for property damage, personal injury, or economic loss caused by the product.

Characterization

To ensure consistently high-quality reagents, each batch of monoclonal antibody is tested for conformance with characteristics of a standard reagent. Representative flow cytometric data is included in this data sheet.

Warning All products contain sodiumazide. This chemical is poisonous and hazardous. Handling should be done by trained staff only.

Explanation of used symbols

	Consult instructions for use
	Catalogue number
	Sufficient for
	In Vitro Diagnostic medical device
	Caution, consult accompanying document
	Keep away from (sun)light
	Biological risks
	Temperature limitation (°C)
	For Research Use Only
	Batch code
	Use by yyyy-mm-dd
	Manufacturer
	Authorized Representative in the European Community
	Conformité Européenne (European Conformity)

		Label - tandem	Ex -max (nm)	Em -max (nm)
P	PURE	purified material	-	-
F	FITC	FITC	488	519
R	R-PE	PE	488, 532	578
C	CyQ	PE-Cy5.18	488, 532	667
A	APC		595, 633, 635, 647	660
PC	PerCP		488, 532	678
PCC	PerCP-Cy5.5		488, 532	695



IQ Products BV
Rozenburglaan 13a
9727 DL Groningen, The Netherlands
 +31 (0)50 57 57 000
 +31 (0)50 57 57 002
 Technical marketing@iqproducts.nl
 Orders orders@iqproducts.nl
 www.iqproducts.nl

 **Products**
bright fluorescence