

## PRODUCT INFORMATION SHEET

### Monoclonal antibodies detecting human antigens

**CD22** FITC    **CD56** R-PE    **RUO**    **REF**    IQP-244FR    50 tests

**RUO**    *For Research Use Only*



#### Description

CD22 antibodies are used as a pan B cell reagent, for the immunophenotyping of B cell lymphomas and HCL. It is more strongly expressed on prolymphocytic leukemia and HCL than in chronic lymphocytic leukemia. B cell lineage ALL, express membrane and cytoplasmic CD22. CD56 is applied for the detection of NK cells in immune monitoring. NK cells form a distinct subpopulation of lymphocytes, which are capable of performing their cytotoxic activity without MHC restriction. Target cells may include virally infected cells or transformed cells and CD56 appears to be involved in the cytotoxic activity of NK cells.

#### CD22

**Clone** B-ly8

**Isotype** Murine IgG1

**Specificity** CD22 forms a loose complex with the BcR cell antigen receptor (BcR). The cytoplasmic domain is tyrosine phosphorylated upon ligation of the BCR and associates via SH2 domains with the tyrosine phosphatase SHP-1, the tyrosine kinase Syk and phospholipase C-g1. CD22 down-modulates the B cell activation threshold, presumably through its association with SHP-1 and other signaling molecules. Mice deficient in CD22 show exaggerated antibody responses to antigen and have raised levels of autoantibodies. CD22 can also mediate cell adhesion through its interaction with cell surface molecules bearing the appropriate sialoglyco-conjugates, but only when these conjugates are not on the CD22 bearing cell itself.

#### HLDA Workshop

The Leukocyte Antigen Factsbook - Barclay, A.J. et.al., Academic Press. London (1997)

#### CD56

**Clone** MOC-1

**Isotype** Murine IgG1

**Specificity** Certain subtypes of T cell lymphomas may express the CD56 antigen. These include peripheral T cell lymphoma (NK cell lymphoma: CD56+, EBV+) and T lymphocytic lymphoma (CD56+/-). In addition, the neoplastic counterpart of the NK cell can be characterized as T gamma lymphocytosis, showing expression of a number of antigens, e.g. CD56+, CD7+, CD16+, CD2+/- and CD8+/. Co-expression of CD56 and CD138 (monoclonal antibody B-A38) is an indication of plasma cell malignancy in multiple myeloma, although it does not occur in all samples.

#### HLDA Workshop


6<sup>th</sup> International Workshop on Human Leukocyte Differentiation Antigens

#### Usage

All these reagents are effectively formulated for direct immunofluorescent staining of human tissue for flow cytometric analysis using 10 µl/10<sup>6</sup> leukocytes for singles and 20 µl/10<sup>6</sup> leukocytes in case of dual and triple combinations. Since applications vary, each investigator should titrate the reagent to obtain optimal results.

#### Representative Data

IQP-244FR (CD22/CD56) was analyzed by flow cytometry using a blood sample from a healthy volunteer. Direct staining was performed by adding 20 µl of this dual to 100 µl blood sample.



## Immunofluorescence staining and lysing protocol

### Flow cytometry method for use with dual and triple combinations

1. Add 100 µl of EDTA-treated blood (i.e. approx.  $10^6$  leukocytes) to a 5 ml reagent tube. The content of one tube is sufficient to perform one test.  
**For combinations with anti-kappa and/or anti-lambda Ig see application note below.**
2. Add to each tube 20 µl of labeled monoclonal antibody combination\*.
3. Vortex the tube to ensure thorough mixing of antibody and cells.
4. Incubate the tube for 15 minutes at room temperature in the dark.
5. Add 100 µl of IQ Lyse (IQP-199 ready-to-use) and mix immediately.
6. Incubate for 10 minutes at room temperature in the dark.
7. Add 2 ml of demineralized water and incubate for 10 minutes in the dark.
8. Centrifuge the labeled cell suspension for 2 minutes at 1000 x g.
9. Remove the supernatant and resuspend the cells in 200 µl of PBS\*\*.
10. Analyze by flow cytometry within four hours (alternatively, the cells may be fixed by 0.05% of formaline in buffered saline for analysis the next day. Some antigens are readily destroyed upon fixation and this should be taken into account when using this alternative).

\* Appropriate mouse Ig isotype control samples can be included in any labeling study

\*\* PBS: Phosphate Buffered Saline, pH 7.2

### **Application note for anti-kappa and/or anti-lambda Ig combinations**

Add 2 ml of PBS containing 0.001% (v/v) Heparin (**prewarmed to 37 °C**) to the cell suspension

Vortex, centrifuge (2 min at 300x g) and discard the supernatant

Repeat this step twice

Resuspend the pelleted blood cells in 100 µl PBS containing 0.001% (v/v) Heparin



### **Handling and Storage**

Antibodies are supplied either as 100 tests per vial (1 ml) for singles or 50 tests per vial (1 ml) for dual and triple combinations. They are supplied in 0.01 M sodium phosphate, 0.15 M NaCl; pH 7.3, 0.2% BSA, 0.09% sodiumazide (NaN<sub>3</sub>). Store the vials at 2-8 °C. Monoclonal antibodies should be protected from prolonged exposure to light. Reagents are stable for the period shown on the vial label when stored properly.

### **Warranty**

Products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied, which extend beyond the description on the label of the product. IQ Products is not liable for property damage, personal injury, or economic loss caused by the product.

### **Characterization**

To ensure consistently high-quality reagents, each batch of monoclonal antibody is tested for conformance with characteristics of a standard reagent. Representative flow cytometric data is included in this data sheet.

### **Warning**

All products contain sodiumazide. This chemical is poisonous and hazardous. Handling should be done by trained staff only.

### Explanation of used symbols



Consult instructions for use



Catalogue number



Sufficient for



In Vitro Diagnostic medical device



Caution, consult accompanying document



Keep away from (sun)light



Biological risks



Temperature limitation (°C)



For Research Use Only



Batch code



Use by yyyy-mm-dd



Manufacturer



Authorized Representative in the European Community



Conformité Européenne (European Conformity)

		<b>Label - tandem</b>	<b>Ex -max (nm)</b>	<b>Em -max (nm)</b>
P	PURE	purified material	-	-
F	FITC	FITC	488	519
R	R-PE	PE	488, 532	578
C	CyQ	PE-Cy5.18	488, 532	667
A	APC		595, 633, 635, 647	660
PC	PerCP		488, 532	678
PCC	PerCP-Cy5.5		488, 532	695



IQ Products BV

Rozenburglaan 13a  
9727 DL Groningen, The Netherlands

☎ +31 (0)50 57 57 000

✉ +31 (0)50 57 57 002

✉ Technical [marketing@iqproducts.nl](mailto:marketing@iqproducts.nl)

✉ Orders [orders@iqproducts.nl](mailto:orders@iqproducts.nl)

✉ [www.iqproducts.nl](http://www.iqproducts.nl)

**bright fluorescence**